


**MINISTÉRIO DA EDUCAÇÃO
UNIVERSIDADE FEDERAL RURAL DA AMAZÔNIA - UFRA
PROGRAMA DE PÓS-GRADUAÇÃO EM AGRONOMIA**

MONYCK JEANE DOS SANTOS LOPES

**BIOPROMOTORES DE CRESCIMENTO EM *Brachiaria brizantha* cv. BRS PIATÃ:
SELEÇÃO, ASPECTOS MORFOFISIOLÓGICOS, ANATÔMICOS E
BIOQUÍMICOS**

**BELÉM
2017**

NC 54796
NOB 53032



MONYCK JEANE DOS SANTOS LOPES

**BIOPROMOTORES DE CRESCIMENTO EM *Brachiaria brizantha* cv. BRS PIATÃ:
SELEÇÃO, ASPECTOS MORFOFISIOLÓGICOS, ANATÔMICOS E
BIOQUÍMICOS**

Tese apresentada ao Programa de Pós-Graduação
em Agronomia da Universidade Federal de Rural
da Amazônia, como requisito para obtenção do
título de Doutor em Agronomia.

Orientadora: Profª. Dra. Gisele Barata da Silva

Co-orientador: PhD. Moacyr Bernardino Dias-
Filho

Biblioteca



44300019

**BELÉM
2017**

TESE
63208
L 864
EX. 01

UNIVERSIDADE FEDERAL DO RIO DE JANEIRO
BIBLIOTECA
F 127.4 - 133.18

Universidade Federal Rural da Amazônia
BIBLIOTECA
Nº 44300019, de 21/12/19

Lopes, Monyck Jeane dos Santos

Biopromotores de crescimento em *Brachiaria brizantha* cv. BRS
Piatã: seleção, aspectos morfofisiológicos, anatômicos e bioquímicos /
Monyck Jeane dos Santos Lopes. – Belém, 2017.
76 f.

Tese (Doutorado em Agronomia) – Programa de Pós-graduação
em Agronomia, Universidade Federal Rural da Amazônia, Belém,
2017.

Orientador: profa. Dra. Gisele Barata da Silva.

Co-orientador: prof. PhD Moacyr Bernardino Dias-filho.

1. *Brachiaria* (Syn. *Urochloa*) *brizantha* cv. Piatã 2. Método de
inoculação – Microrganismos 3. Crescimento - Gramínea forrageira 4.
Aspectos - Morfofisiológicos – Anatômicos – Bioquímicos I. Silva,
Gisele Barata da, (orient.) II. Dias-filho, Moacyr Bernardino, (co-
orient.) III. Título.

CDD – 633.208



**MINISTÉRIO DA EDUCAÇÃO
UNIVERSIDADE FEDERAL RURAL DA AMAZÔNIA
PROGRAMA DE PÓS-GRADUAÇÃO EM AGRONOMIA**

MONYCK JEANE DOS SANTOS LOPES

**BIOPROMOTORES DE CRESCIMENTO EM *Brachiaria brizantha* cv. BRS
PIATÁ: SELEÇÃO, ASPECTOS MORFOFISIOLÓGICOS, ANATÔMICOS E
BIOQUÍMICOS**

Tese apresentada ao Programa de Pós-Graduação em Agronomia, da Universidade
Federal Rural da Amazônia, como requisito para obtenção do título de Doutor em
Agronomia.

20 de junho de 2017

BANCA EXAMINADORA

Profª. Dra. Gisele Barata da Silva – Presidente
UNIVERSIDADE FEDERAL RURAL DA AMAZÔNIA

Dra. Fernanda Ilki Borges de Souza – 1º Examinador
EMBRAPA AMAZÔNIA ORIENTAL

Dr. Marcelo Murad Magalhães – 2º Examinador
EMBRAPA AMAZÔNIA ORIENTAL

Dr. Fábio Lopes Olivares – 3º Examinador
UNIVERSIDADE ESTADUAL DO NORTE FLUMINENSE

Dr. Walter Vellascó Duarte Silvestre – 4º Examinador
UNIVERSIDADE FEDERAL RURAL DA AMAZÔNIA

Dr. Roberto Lisboa Cunha (Suplente) – 5º Examinador
EMBRAPA AMAZÔNIA ORIENTAL

AGRADECIMENTOS

Ao grandioso Jeová Deus e ao seu filho Jesus Cristo por todas as maravilhosas bênçãos que me concedem diariamente.

À Universidade Federal Rural da Amazônia (UFRA) pela oportunidade de realizar o curso de pós-graduação em Agronomia;

À Fundação Amazônia de Amparo a Estudos e Pesquisas - FAPESPA, pela concessão da bolsa de estudos;

A minha orientadora Dra. Gisele Barata, pelo apoio, ensinamento e compreensão;

Aos membros da banca, pela sua disponibilidade a compartilhar seu conhecimento;

Ao Dr. Eniel David Cruz, Dr. Walter Vellasco, M.Sc. Lenilson Silva e Sr. Zezinho pelo suporte técnico;

A todos os meus colegas do Laboratório de Proteção de Plantas (LPP);

Aos meus familiares e amigos, pelo incentivo e apoio;

A três anjos que Deus colocou na minha vida, que sem eles esse título seria impossível:

- Meu amado esposo Moacyr Dias-Filho, por todo amor, compreensão, positividade e ajuda em todos os momentos.

- A minha amiga Bruna Calzavara, uma mãe de coração, que me ajudou imensamente com meu filho, para que eu pudesse me dedicar ao doutorado.

- Ao graduando Thomaz Castro, um estagiário que me socorreu nos experimentos, muitas vezes dando continuidade sozinho, quando minha saúde não permitia.

Ao meu presente mais precioso, meu amado filho Moacyr Dias-Neto, que me deu forças a cada momento e em cada dificuldade.

Enfim, a todos que direta e/ou indiretamente me apoiaram para o término dessa tese.

Meus sinceros agradecimentos!

LISTA DE FIGURAS

ARTIGO 1

Fig. 1 Shoot (a-e) and root (f-j) of *Brachiaria brizantha*, 21 days after inoculation (35 days after seedling emergence). Unfertilized-control (a, f), fertilized-control (b, g), inoculated with *Pseudomonas fluorescens* (BRM-32111) (c, h), *Burkholderia pyrrocinia* (BRM-32113) (d, i) and co-inoculated with BRM-32111 + BRM-32113 (e, j).344

Fig. 2 Number of leaves per plant (a), plant height (H) (b), culm length (c) and SPAD index (d) of *Brachiaria brizantha* inoculated with growth-promoting rhizobacteria. Values are means $\pm$ SE ($n = 5$). Days = days after sowing. C - = unfertilized-control; C + = fertilized-control; B = *Burkholderia pyrrocinia*; P = *Pseudomonas fluorescens*; MIX = B + P.35

Fig. 3 *Brachiaria brizantha* responses to growth-promoting rhizobacteria. (a) Net photosynthesis (A), (b) stomatal conductance (gs) and (c) transpiration (E). At 21 days after inoculation (35 days after seedling emergence). Columns with different letters are significantly different among treatments ($P < 0.05$, Duncan Test). C - = unfertilized-control; C + = fertilized-control; B = *Burkholderia pyrrocinia*; P = *Pseudomonas fluorescens*; MIX = B + P.36

Fig. 4 *Brachiaria brizantha* responses to growth-promoting rhizobacteria. (a) Shoot dry mass production (SDM), (b) root dry mass production (RDM), (c) total dry mass production (TDM) and, (d) biomass allocation pattern. 21 days after inoculation (35 days after seedling emergence). Columns with different letters are significantly different among treatments ($P < 0.05$, Duncan Test). Different upper-case letters within columns indicate significant differences among plant organs ($P < 0.05$, Duncan Test). C - = unfertilized-control; C + = fertilized-control; B = *Burkholderia pyrrocinia*; P = *Pseudomonas fluorescens*; MIX = B + P.37

Fig. 5 *Brachiaria brizantha* responses to growth-promoting rhizobacteria. (a) Free ammonium, (b) nitrate, (c) amino acid and, (d) total soluble proteins. 21 days after inoculation (35 days after seedling emergence). Columns with different letters are significantly different among treatments ($P < 0.05$, Duncan Test). C - = unfertilized-control; C + = fertilized-control; Mix = *Burkholderia pyrrocinia* + *Pseudomonas fluorescens*.38

Fig. 6 Nitrogen concentration of *Brachiaria brizantha* inoculated with growth-promoting rhizobacteria. 21 days after inoculation (35 days after seedling emergence). Columns with different letters are significantly different among treatments ($P < 0.05$, Duncan Test). C - = unfertilized-control; C + = fertilized-control; Mix = *Burkholderia pyrrocinia* + *Pseudomonas fluorescens*.38

ARTIGO 2

Figura 1 Secção transversal de folha (a-c), colmo (d-f) e raiz (g-i) de *Brachiaria brizantha* não inoculada sem fertilizante (a, d, g), com fertilizante (b, e, h) e coinoculada com *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113). 21 dias após a inoculação (35 dias após a emergência das plantas). Célula buliforme (Bc), bainha do feixe (Bs), feixe vascular (Vb), xilema (X), mesófilo (Mes), exoderme (Ex), periciclo (pe), cilindro vascular (vc) e protoxilema (px). 55

Figura 2 Clorofila (a), índice Spad (b), fotossíntese líquida (A) (c) e eficiência do uso da água (WUE) de *Brachiaria brizantha* não inoculada sem fertilizante (C-), com fertilizante (C+) e coinoculada com *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113) (Mix). Médias seguidas da mesma letra na coluna não diferem ($P < 0,05$, teste de Duncan).56

Figura 3 Amido (a) e carboidratos totais (b) de *Brachiaria brizantha* não inoculada sem fertilizante (C-), com fertilizante (C+) e coinoculada com *Pseudomonas fluorescens* (BRM-

32111) e *Burkholderia pyrrocinia* (BRM-32113) (Mix). Médias seguidas da mesma letra em cada coluna não diferem ($P < 0,05$, teste de Duncan)
566

Figura 4 Altura (H) (a), área radicular (b) e produção de biomassa (c) de *Brachiaria brizantha* não inoculada sem fertilizante (C-), com fertilizante (C+) e coinoculada com *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113) (Mix). Médias seguidas da mesma letra na coluna não diferem ($P < 0,05$, teste de Duncan)..... 57

Figura 5 *Brachiaria brizantha* coinoculada com rizobactérias promotoras de crescimento em plantas (PGPR), 21 dias após a inoculação (35 dias após a emergência das plantas). *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113) modificaram a anatomia (folha, colmo e raiz), incrementaram a altura (H), taxa fotossintética (A), eficiência do uso da água (WUE), concentração de clorofila, N, P, Na, Mg, Fe, amido, carboidrato, proteína, área radicular, aumentando a qualidade de forragem e produção de biomassa em *B. brizantha*..... 58

ARTIGO 3

Figure 1 *Brachiaria brizantha* cv. Piatã under full-sun (a-d) and shade (e-h). 21 days after inoculation (35 days after seedling emergence). Non-inoculated plant (a, e), inoculated with *Pseudomonas fluorescens* (BRM-32111) (b, f), inoculated with *Burkholderia pyrrocinia* (BRM-32113) (c, g) and co-inoculated with BRM-32111 + BRM-32113 (d, h)..... 73

Figure 2 Light intensity and plant growth-promoting rhizobacteria (PGPR) on plant height (H) of *Brachiaria brizantha*. Columns with different letters are significantly different among PGPR inoculation ($P < 0.05$, Duncan's test). Mean $\pm$ s.e. ($n = 5$). C = Non-inoculated; BRM-32111 = *Pseudomonas fluorescens*; BRM-32113 = *Burkholderia pyrrocinia*; MIX = BRM-32111 + BRM-32113..... 73

Figure 3 Light intensity and plant growth-promoting rhizobacteria (PGPR) on specific leaf area (SLA) and leaf area ratio (LAR) of *Brachiaria brizantha*. Columns with different letters are significantly different among PGPR inoculation ($P < 0.05$, Duncan's test). Mean $\pm$ s.e. ($n = 5$). C = Non-inoculated; BRM-32111 = *Pseudomonas fluorescens*; BRM-32113 = *Burkholderia pyrrocinia*; MIX = BRM-32111 + BRM-32113. 74

Figure 4 Light intensity and plant growth-promoting rhizobacteria (PGPR) on leaf area (LA) and SPAD index (chlorophyll content) of *Brachiaria brizantha*. Columns with different letters are significantly different among PGPR inoculation ($P < 0.05$, Duncan's test). Mean $\pm$ s.e. ($n = 5$). C = Non-inoculated; BRM-32111 = *Pseudomonas fluorescens*; BRM-32113 = *Burkholderia pyrrocinia*; MIX = BRM-32111 + BRM-32113 74

Figure 5 Light intensity and plant growth-promoting rhizobacteria (PGPR) on total dry mass production (TDM) and relative growth rate (RGR) of *Brachiaria brizantha*. Columns with different letters are significantly different among PGPR inoculation ($P < 0.05$, Duncan's test). Mean $\pm$ s.e. ($n = 5$). C = Non-inoculated; BRM-32111 = *Pseudomonas fluorescens*; BRM-32113 = *Burkholderia pyrrocinia*; MIX = BRM-32111 + BRM-32113..... 75

Figure 6 Light intensity and plant growth-promoting rhizobacteria (PGPR) on biomass allocation pattern of *Brachiaria brizantha*. Columns with different lower case letters are significantly different among treatments ($P < 0.05$, Duncan Test). Different upper-case letters within columns indicate significant differences among plant organs ($P < 0.05$, Duncan's test). Mean $\pm$ s.e. ($n = 5$). BRM-32111 = *Pseudomonas fluorescens*; BRM-32113 = *Burkholderia pyrrocinia*; MIX = BRM-32111 + BRM-32113. 75

LISTA DE TABELAS

ARTIGO 1

Table 1 Inoculation method of plant growth-promoting rhizobacteria on the biomass production of <i>Brachiaria brizantha</i> . Shoot dry mass (SDM), root dry mass (RDM), total dry mass (TDM).	32
Table 2 Effects of fertilization and plant growth-promoting rhizobacteria on the biomass production of <i>Brachiaria brizantha</i> . Shoot dry mass (SDM), root dry mass (RDM), total dry mass (TDM).	32
Table 3 Leaf area (LA cm ²), root dry mass (RDM - g day ⁻¹) /shoot dry mass (SDM) ratio, relative growth rate (RGR), number of tillers (NT), expanded leaf length (EF - cm), leaf appearance rate (LApR - L ⁻¹ day ⁻¹) and leaf elongation rate (LER - cm day ⁻¹) of <i>Brachiaria brizantha</i> with growth-promoting rhizobacteria.	33
Table 4 Correlation coefficient (<i>r</i>) of the correlations between SPAD, leaf area (LA), plant height (H), number of tillers (NT), photosynthesis (A) and nitrogen concentration (N) versus total dry mass (TDM) of <i>Brachiaria brizantha</i> .	33

ARTIGO 2

Tabela 1 Parâmetros anatômicos (μm or μm ²) de <i>Brachiaria brizantha</i> não inoculada sem fertilizante (C-), com fertilizante (C+) e coinoculada com <i>Pseudomonas fluorescens</i> (BRM-32111) e <i>Burkholderia pyrrocinia</i> (BRM-32113) (Mix).	53
Tabela 2 Concentração de macro e micronutrients (g/kg), fibra em detergente neutro (FDN - g kg ⁻¹ of DM), fibra em detergente ácido (FDA - g kg ⁻¹ of DM) e proteína bruta (PB - g kg ⁻¹ of DM) de <i>Brachiaria brizantha</i> não inoculada sem fertilizante (C-), com fertilizante (C+) e coinoculada com <i>Pseudomonas fluorescens</i> (BRM-32111) e <i>Burkholderia pyrrocinia</i> (BRM-32113) (Mix).	54

ARTIGO 3

Table 1 Light intensity and plant growth-promoting rhizobacteria (PGPR) on number of tillers (NT), number of leaves per plant (NL), expanded leaf length (EF - cm), culm length (C - cm), leaf appearance rate (LApR - L ⁻¹ .day ⁻¹), leaf elongation rate (LER - L ⁻¹ .day ⁻¹) and culm elongation rate (CER - L ⁻¹ .day ⁻¹) of <i>Brachiaria brizantha</i> .	72
Table 2 Light intensity and plant growth-promoting rhizobacteria (PGPR) on the leaf (LDM - g), culm (CDM - g), root (RDM - g) and root/shoot dry mass production, and allometric coefficient K (growth rate of root and shoot ratio) of <i>Brachiaria brizantha</i> .	72

LISTA DE ABREVIATURAS

BRM-32111 - *Pseudomonas fluorescens*

BRM-32113 - *Burkholderia pyrrocinia*

MIX - *Pseudomonas fluorescens* + *Burkholderia pyrrocinia*

PGPR – Rizobactérias promotoras de crescimento em plantas

H - Altura

TDM - Massa seca total

SDM - Massa seca da parte aérea

RDM – Massa seca radicular

RGR - Taxa de crescimento relativo

g_s - Condutância estomática

E - Transpiração

A - Fotossíntese

WUE – Eficiência do uso de água

LA - Área foliar

SLA - Área foliar específica

LApR - Taxa de aparecimento foliar

LER - Taxa de alongamento foliar

CER - Taxa de alongamento do colmo

NT - Número de perfilhos

EF- Folha expandida

NL – Número de folhas

NDF – Fibra detergente neutro

ADF – Fibra detergente ácido

RESUMO

O uso de microrganismos benéficos em pastagens é uma técnica potencialmente vantajosa para o manejo sustentável, reduzindo a adubação química na manutenção das pastagens ainda produtivas e na recuperação de áreas degradadas, em sistemas convencionais e integrados. Os objetivos deste estudo foram: determinar o melhor método de inoculação de microrganismos capazes de promover o crescimento da gramínea forrageira *Brachiaria* (Syn. *Urochloa*) *brizantha* cv. Piatã; examinar o seu efeito sobre aspectos morfológicos, anatômicos, fisiológicos, bioquímicos e nutricionais; e verificar o efeito dos biopromotores no crescimento dessa gramínea sob diferentes intensidades luminosas. A seleção foi *in vivo* com três métodos de inoculação: sementes, sementes e rega, e apenas por rega do solo. Os ensaios foram conduzidos em casa de vegetação e em viveiro sob pleno sol e sombra na Universidade Federal Rural da Amazônia (UFRA) (01°27'25"S, 48°26'36"W), em Belém, Pará, Brasil. Foram avaliados parâmetros de morfogênese, anatômicos, fisiológicos, bioquímicos e nutricionais. O uso de *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113), coinoculadas pela rega do solo, modificou a anatomia (folha, colmo e raiz), incrementou as taxas fotossintéticas (14%), eficiência do uso da água (22%), índice Spad (37%), clorofila (122%), nitrato (100%), proteínas (80%), carboidratos totais (29%), amido (133%), N (30%), P (20%), Mg (20%) e Fe (180%), promovendo uma maior qualidade e produção de biomassa em *B. brizantha*. Também promoveram o crescimento em *B. brizantha* sob diferentes condições luminosas, incrementando em mais de 100% a produção de biomassa, em comparação as plantas controles com e sem adubo. O método de inoculação e a intensidade de luz foram fundamentais para determinar o potencial do microrganismo como promotor do crescimento em *B. brizantha*. Nosso estudo evidencia o potencial biofertilizante da coinoculação de *P. fluorescens* e *B. pyrrocinia* em *B. brizantha* cv. BRS Piatã sob diferentes intensidades de luz, possibilitando seu uso em sistemas convencionais e integrados, contribuindo para a obtenção de pastagens mais produtivas e de maior qualidade.

Palavras-chave: Inoculação. Forrageira. BRS Piatã. *Pseudomonas fluorescens*. *Burkholderia pyrrocinia*. Biomassa.

ABSTRACT

The inoculation of beneficial microorganisms in forage grasses is a potentially advantageous technique for sustainable pasture management by decreasing the need for chemical fertilization to maintain pasture productivity and reclaim degraded pastures, under both conventional and integrated systems (i.e., subjected to shade). The aim of this study was to determine the most effective method of microorganism inoculation on *Brachiaria* (Syn. *Urochloa*) *brizantha* cv. BRS Piatã, by examining its effect on selected morphological, anatomical, physiological, biochemical and qualitative traits. In addition, we examined the impact of inoculation on the growth characteristics of this forage grass cultivar, grown under contrasting light conditions. Screening tests were carried out *in vivo*, for three methods of inoculation: in the seed, seed and soil, and soil. The experiment was conducted in a greenhouse, under full sun and shade at the Federal Rural University of Amazonia (UFRA) (01°27'25 "S, 48°26'36" W) in Belém, Pará, Brazil. *Pseudomonas fluorescens* (BRM-32111) and *Burkholderia pyrrocinia* (BRM-32113), when co-inoculated by soil drenching, modified the anatomy (leaf, stem, root), increased the photosynthetic rate (14%), water use efficiency (22%), Spad index (37%), chlorophyll (122%), nitrate (100%), proteins (80%), total carbohydrates (29%), starch (133%), N (30%), P (20%), Mg (20%) and Fe (180%) contents and increased forage quality and biomass production of *B. brizantha* cv. BRS Piatã. Co-inoculation promoted growth of *B. brizantha* under contrasting light conditions, increasing biomass production by more than 100% relative to unfertilized- and fertilized-control (non-inoculated) plants. Our results imply that the method of inoculation and the light environment are important in determining the potential of microorganism for promoting plant growth. This study attests that co-inoculation of *P. fluorescens* and *B. pyrrocinia* are effective biofertilizers to increase *B. brizantha* productivity and forage quality under full sun and shaded environments.

Keywords: Inoculation. Forage grass. BRS Piatã. *Pseudomonas fluorescens*, *Burkholderia pyrrocinia*. Biomass.

SUMÁRIO

RESUMO.....	10
ABSTRACT	11
1. CONTEXTUALIZAÇÃO.....	13
REFERÊNCIAS	15
2. <i>Pseudomonas fluorescens</i> AND <i>Burkholderia pyrrocinia</i> GROWTH IMPROVEMENT AND PHYSIOLOGICAL RESPONSES OF <i>Brachiaria brizantha</i>	17
Abstract	17
2.1 Introduction	18
2.2 Materials and methods.....	20
2.3 Results.....	23
2.4 Discussion	25
REFERENCES	30
Tables.....	32
Figures	34
3. VALOR NUTRITIVO, PRODUÇÃO DE BIOMASSA E PARÂMETROS ANATÔMICOS DE <i>Brachiaria brizantha</i> COM RIZOBACTÉRIAS PROMOTORAS DE CRESCIMENTO (PGPR)	39
RESUMO.....	39
3.1 Introdução	40
3.2 Material e métodos	41
3.3 Resultados	43
3.4 Discussão	45
REFERÊNCIAS	49
Tabelas.....	53
Figuras	55
4. LIGHT AND PLANT GROWTH-PROMOTING RHIZOBACTERIA (PGPR) EFFECTS ON <i>Brachiaria brizantha</i> GROWTH AND PHENOTYPIC PLASTICITY TO SHADE	59
4.1 Abstract	59
4.2 Introduction	60
4.3 Materials and methods.....	61
4.4 Results.....	64
4.5 Discussion	66
REFERENCES	69
Tables.....	72
Figures	73
CONCLUSÕES GERAIS	76

CONTEXTUALIZAÇÃO

O Brasil é um dos maiores produtores e exportadores de carne bovina do mercado mundial (USDA, 2017). A razão para isso é que a criação do rebanho bovino brasileiro é a pasto, o que torna a produção mais econômica. As pastagens brasileiras tem grande potencial produtivo, mas apenas 30% desse potencial esta sendo explorado (STRASSBURG et al., 2014). Uma das principais causas dessa baixa produtividade é o aumento de pastagens degradadas (DIAS-FILHO, 2011).

Para atender a crescente demanda mundial por produtos de origem animal, com menor impacto ambiental, é necessário aumentar a produtividade das pastagens ainda ativas e recuperar as que estão degradadas (DIAS-FILHO, 2011; PAGANO et al., 2017). Uma das formas de aumentar a produção nos pastos do Brasil, sem maior expansão das áreas, é pelo manejo da fertilidade do solo, mas seu uso é limitado, pois as adubações periódicas têm alto custo (DIAS-FILHO, 2011). Uma solução para uma produção menos onerosa e com menor impacto ambiental seria substituir os insumos químicos por biopromotores de crescimento do vegetal (PAGANO et al., 2017; TIMMUSK et al., 2017; DUCHENE et al., 2017).

Os biopromotores são microrganismos benéficos que promovem o crescimento vegetal por regular a rota metabólica dos fitormônios e tornar os nutrientes do solo disponíveis (DOORNBOS et al., 2012; PAGANO et al., 2017; TIMMUSK et al. 2017). O efeito positivo das rizobactérias promotoras de crescimento é relatado em várias Poaceae, como: arroz (REGAR & YADAV, 2017), cana-de-açúcar (SANTOS et al., 2017), milho (CALVO et al., 2017) e forrageiras (HUNGRIA et al., 2016; MARQUES et al., 2017).

No Brasil, cerca de 80% das áreas de pastagens são formadas por gramíneas do gênero *Brachiaria* (Syn. *Urochloa*), uma forrageira perene, com elevado crescimento, tolerância ao pastejo e alta disponibilidade de forragem (REIS et al., 2013; PAGANO et al., 2017). Dentre as cultivares de *B. brizantha* disponíveis no mercado, destaca-se a BRS Piatã, por apresentar menor alongamento da haste e sazonalidade na produção, maior acúmulo de forragem durante o período seco e desempenho animal, e ser mais tolerante à cigarrinha-das-pastagens em relação as outras cultivares dessa espécie (REIS et al., 2013).

No entanto, o alto potencial alelopático de *Brachiaria* (SOUZA FILHO et al., 2005; KATO-NOGUCHI et al., 2014) e as diferentes intensidades de luz nas pastagens em sistemas convencionais e integrados, podem interferir na interação mutualística entre plantas e

microrganismos benéficos (AGUILAR-CHAMA & GUEVARA, 2016; KONVALINKOVÁ & JANSÁ, 2016; VIMAL et al., 2017).

Estudos prévios com *Pseudomonas* sp. e *Burkholderia* sp. atestam seu potencial como promotor de crescimento em arroz (RÊGO et al., 2014, NASCENTE et al., 2016). Estas rizobactérias são conhecidas por aumentarem a síntese de auxina, absorção de nutrientes, teor de clorofila, taxa fotossintética e produção de biomassa (AHMAD & KIBRET, 2014). Nossa hipótese é que *Burkholderia pyrrocinia* e *Pseudomonas fluorescens* também são capazes de promover o crescimento em *Brachiaria brizantha* cv. BRS Piatã (Hochst. ex A. Rich.) Stapf., aumentando a produtividade das pastagens, em sistemas convencionais e integrados, de uma forma sustentável e promissora. Portanto, nosso objetivo geral é determinar o melhor método de inoculação de *B. pyrrocinia* e *P. fluorescens* e verificar seus efeitos no crescimento em *B. brizantha* cv. BRS Piatã. Para alcançar o objetivo, essa pesquisa foi dividida nos seguintes artigos:

Artigo 1: *Pseudomonas fluorescens* and *Burkholderia pyrrocinia* growth improvement and physiological responses of *Brachiaria brizantha*

Esse artigo apresenta o melhor método de inoculação de *P. fluorescens* e *B. pyrrocinia*, verificando a sua influência sobre as características morfológicas, fisiológicas e bioquímicas de *B. brizantha*.

Artigo 2: Valor nutritivo, produção de biomassa e parâmetros anatômicos de *Brachiaria brizantha* com rizobactérias promotoras de crescimento (PGPR)

Nesse artigo avaliam-se os efeitos da co-inoculação de *P. fluorescens* e *B. pyrrocinia* na anatomia, bioquímica, qualidade nutricional e produção de biomassa de *B. brizantha*.

Artigo 3: Light and plant growth-promoting rhizobacteria (PGPR) effects on *Brachiaria brizantha* growth and phenotypic plasticity to shade

Nesse artigo avaliam-se os efeitos da intensidade de luz e da inoculação de *P. fluorescens* e *B. pyrrocinia* no crescimento, plasticidade fenotípica e tolerância à sombra de *B. brizantha*.

REFERÊNCIAS

- AGUILAR-CHAMA, A., GUEVARA, R. Resource allocation in an annual herb: Effects of light, mycorrhizal fungi, and defoliation. **Acta Oecol.** v.71, p.1-7, 2016.
- AHEMAD M. & KIBRET M. Mechanisms and applications of plant growth promoting rhizobacteria: Current perspective. **JKSU– Science** v.26, p.1–20, 2014.
- CALVO P., WATTS D.B., KLOEPFER J.W. and TORBERT H.A. Effect of microbial-based inoculants on nutrient concentrations and early root morphology of corn (*Zea mays*) **J. Plant Nutrit and Soil Sci** v.180, p.56–70, 2017.
- DIAS-FILHO, M.B. Degradação de pastagens: processos, causas e estratégias de recuperação. 4. Ed., rev. atual. e ampl. Belém, PA. 2011.
- DOORNBOS, R.F, LOON L.C.V., BAKKER, P.A.H.M. Impact of root exudates and plant defense signaling on bacterial communities in the rhizosphere. A review. **Agron. Sustain. Dev.** v.32, p.227–243, 2012.
- DUCHENE, O., VIAN, J.F., CELETTE F. Intercropping with legume for agroecological cropping systems: Complementarity and facilitation processes and the importance of soil microorganisms. A review. **Agriculture, Ecosystems and Environment**, v.240, p.148–161, 2017.
- HUNGRIA, M., NOGUEIRA, M.A., ARAUJO, R.S. Inoculation of *Brachiaria* spp. with the plant growth-promoting bacterium *Azospirillum brasilense*: An environment-friendly component in the reclamation of degraded pastures in the tropics. **Agr. Ecosyst. Environ.** v.221, p.125–131, 2016.
- KATO-NOGUCHI, H., KOBAYASHI, A., OHNO, O., KIMURA, F., FUJII, Y., SUENAGA, K. Phytotoxic substances with allelopathic activity may be central to the strong invasive potential of *Brachiaria brizantha*. **J. Plant. Physio.** v.171, p.525–530, 2014.
- KONVALINKOVÁ, T. JANSÁ, J. Lights off for arbuscular mycorrhiza: on its symbiotic functioning under light deprivation. **Frontiers in Plant Science**, v.7, 782, 2016,
- MARQUES, A.C.R., OLIVEIRA, L.B., NICOLOSO, F.T., JACQUES, R.J.S., GIACOMINI, S.J., QUADROS, F.L.F. Biological nitrogen fixation in C4 grasses of different growth strategies of South America natural grasslands. **Appl. Soil Ecol.** v. 113, p.54 – 62, 2017.
- NASCENTE, A.S., FILIPPI, M.C.C., LANNA, A.C., SOUZA, A.C.A., LOBO, V.L.S. SILVA, G.B. Biomass, gas exchange, and nutrient contents in upland rice plants affected by application forms of microorganism growth promoters. **Environmental Science and Pollution Research** 2016
- PAGANO M.C., CORREA E.J.A., DUARTE N.F., YELIKBAYEV B., O'DONOVAN A. and GUPTA V.K. Advances in Eco-Efficient Agriculture: The Plant-Soil Mycobiome. **Agriculture**, v.7, n.14, 2017.

REGAR, K.L. and YADAV, J. Influence of PGPR and Zinc Enriched FYM on Growth and Yield of Rice at Different Levels of Phosphors in an Inceptisol of Varanasi, India. **Int.J.Curr.Microbiol.App.Sci** v.6, n.4, p.1453-1464, 2017.

RÊGO, M.C.F., BORGES, F.I., FILIPPI, M.C.C., GONÇALVES, L.A., SILVA, G.B. Morphoanatomical and biochemical changes in the roots of rice plants induced by plant growth-promoting microorganisms. **J. Botany**, 2014.

REIS, A.R., BERNARDES, T.F., SIQUEIRA G.R. Forragicultura: ciência, tecnologia e gestão de recursos forrageiros. 714p. 2013.

SANTOS, S.G., RIBEIRO, F.S., FONSECA, C.S. PEREIRA W., SANTOS L.A., REIS V.M. Development and nitrate reductase activity of sugarcane inoculated with five diazotrophic strains. **Arch Microbiol**, 2017

SOUZA FILHO, A.P.S., PEREIRA, A.A.G., BAYMA, J.C. Aleloquímico produzido pela gramínea forrageira *Brachiaria humidicola*. **Planta Daninha**. v.23, p.25-32, 2005.

STRASSBURG, B.B.N., LATAWIEC, A.E., BARIONI, L.G., NOBRE, C.A., SILVA, V.P., VALENTIM, J.F., VIANNA, M., ASSAD, E.D. 2014. When enough should be enough: improving the use of current agricultural lands could meet production demands and spare natural habitats in Brazil. **Global Environ. Change** v.28, p.84-97, 2014.

TIMMUSK, S., BEHERS, L., MUTHONI, J., MURAYA, A., ARONSSON A.C. Perspectives and challenges of microbial application for crop improvement. **Frontiers in Plant Science**, v.8, article 49, 2017.

USDA, 2017. United States Department of Agriculture. Production, supply and distribution online. <https://apps.fas.usda.gov/psdonline/app/index.html> (accessed 25.05.2017).

VALLE, C.B., BARRIOS, S.C.L., JANK, L., SANTOS, M.F. Melhoramento de plantas forrageiras para uma pecuária de baixa emissão de carbono. In: Pedreira, B.C., Pereira, D.H., Pina, D.S., Carnevalli, R.A., Lopes, L.B. **Intensificação da produção animal em pastagens**. Brasília, DF: Embrapa, pp 109-139. 2014.

VIMAL S.R., SINGH J.S., ARORA N.K. and SINGH S. Soil-Plant-Microbe Interactions in Stressed Agriculture Management: A Review. **Pedosphere**, v.27, p.177-192, 2017.

1 *Pseudomonas fluorescens* and *Burkholderia pyrrocinia* growth improvement and
 2 physiological responses of *Brachiaria brizantha*¹

3 Monyck Jeane dos Santos Lopes, Moacyr Bernardino Dias Filho, Thomaz Henrique dos Reis
 4 Castro, Marta Cristina Corsi de Filippi, Gisele Barata da Silva

5 **ABSTRACT**

6 The use of beneficial microorganisms in forage grasses is a potentially advantageous
 7 technique for a more sustainable pasture management by decreasing the need for chemical
 8 fertilization. Our aims were to determine the best method of microorganism inoculation on
 9 *Brachiaria* (Syn. *Urochloa*) *brizantha* cv. BRS Piatã, compare the responses of inoculated
 10 plants of this forage grass with fertilized and unfertilized controls and examine its effect on
 11 some morphological, physiological and biochemical. On the first experiment, three inoculation
 12 methods were tested: in the seed, seed and soil, and soil, with *Pseudomonas fluorescens* (BRM-
 13 32111) and *Burkholderia pyrrocinia* (BRM-32113). In the second experiment, fertilized and
 14 unfertilized plants were either inoculated with BRM-32111, BRM-32113 and co-inoculated
 15 (BRM-32111 + BRM-32113). In a final experiment, *U. brizantha* was inoculated by soil
 16 drenching with BRM-32111, BRM-32113 and co-inoculated (BRM-32111 + BRM-32113), and
 17 compared to fertilized- and unfertilized-controls. The inoculation by soil drenching, at seedling
 18 stage, was more effective than inoculation only in the seed or both in the seed and by soil
 19 drenching. The fertilizer may have suppressed the beneficial bacterial effects on the growth of
 20 *B. brizantha*. *P. fluorescens* and *B. pyrrocinia* co-inoculated increased nitrate, protein, nitrogen
 21 concentration, Spad index (chlorophyll content), leaf area, number of tillers, net photosynthesis
 22 and total biomass production of *B. brizantha* plants. Our results point out to a potentially

¹ Este capítulo segue as normas de formatação da **Applied Soil Ecology**.

valuable source of practical information in the search of an eco-friendlier approach to increase pasture productivity.

Highlights

- Inoculation method defines the potential of microorganism for promoting plant growth.
- *P. fluorescens* and *B. pyrrocinia* increased *Brachiaria brizantha* biomass production.
- Their co-inoculation was an effective biofertilizer in *B. brizantha*.

Keywords: forage grass, *Urochloa*, plant growth-promoting rhizobacteria, fertilized, nitrate, nitrogen.

1. Introduction

Increasing global population coupled with shifting dietary preferences in emerging economies is leading to a substantial increase in the consumption of livestock products, mainly beef. Sustainable intensification of current pasturelands in developing countries is an important tool to meeting future demands for beef (Strassburg et al., 2014). The use of plant growth-promoting microorganisms in pastures is a potentially advantageous technique, as a more economical and eco-friendlier approach to increase pasture productivity, when compared to the sole use of chemical fertilizers (Dias-Filho, 2011; Kelemu et al., 2011; Hungria et al., 2016; Marques et al., 2017).

Plant growth-promoting microorganisms are beneficial endophytic or rhizospheric microorganisms, able to colonize roots and directly promote growth by regulating the pathway of plant hormones, increasing the biosynthesis of auxin, cytokinin, gibberellin, or minimizing the ACC synthesis, the ethylene precursor, delaying plant senescence (Doornbos et al., 2012; Timmusk et al., 2017). Another beneficial mechanism of growth-promoting microorganisms are to increase the availability of essential nutrients for plant growth, such as nitrogen and phosphorus, and to promote induced resistance of plant defenses against diseases, pests, and abiotic stressors (Doornbos et al., 2012; Timmusk et al., 2017). Growth promotion, resulting

48 from the association of beneficial microorganisms, has been reported in wheat plants (Silveira
49 et al., 2016), forage grass (Marques et al., 2017) and *Sorghum bicolor* (Santos et al., 2017).

50 In Brazil, around 80% of pasture areas are formed by grasses of the *Brachiaria* (Syn.
51 *Urochloa*) genus, among which stands out *B. brizantha*, being the cultivar BRS Piatã one of the
52 major current options for pasture formation (Valle et al., 2014). Most of these pasture areas is
53 under low fertility soils, requiring chemical fertilization to produce satisfactorily and, in
54 particular, nitrogen fertilization to intensify pasture management (Dias-Filho, 2011; Hungria et
55 al., 2016; Marques et al., 2017). However, the efficiency of fertilizer use by plants, particularly
56 that of nitrogen, may vary greatly. This can create adverse environmental impacts by increasing
57 greenhouse gas emissions and eutrophication (Mantelin and Touraine, 2003; Keuter et al., 2014;
58 Pii et al., 2015).

59 The association between *B. brizantha* and diazotrophic bacteria could be highly
60 beneficial, for nitrogen fixation and subsequent transfer of the fixed nitrogen to the host plant,
61 increasing the sustainability of agriculture and reducing its impact on the environment.
62 However, *Brachiaria* genus is known to be very allelopathic (Souza Filho et al., 2005; Kato-
63 Noguchi et al., 2014). This may deter the establishment of beneficial microorganisms in the
64 rhizosphere (Doornbos et al., 2012). Therefore, research on the optimal inoculation method of
65 beneficial microorganisms is crucial for this grass genus.

66 Studies conducted at the Federal Rural University of Amazon, proved that diazotrophic
67 rhizobacteria identified as *Pseudomonas fluorescens* and *Burkholderia pyrrocinia*, isolated
68 from the rhizosphere soil, in Pará, Brazil, are growth promoters in rice plants (Rego et al., 2014;
69 Nascente et al., 2016). We hypothesize that these growth-promoting microorganisms are
70 capable of stimulating growth in *B. brizantha* cv. BRS Piatã, grown in low-fertility soils, being
71 an important strategy for the sustainable intensification of pasture production systems.
72 Therefore, the aim of this study was to determine the optimal inoculation method of plant

73 growth promoting microorganisms, and examine their effect on some morphological,
74 physiological, and biochemical responses of *B. brizantha* cv. BRS Piatã.

75 2. Materials and methods

76 2.1 Study site, plant and soil

77 The experiment was conducted at the Plant Protection Laboratory and greenhouse of
78 the Federal Rural University of Amazonia (UFRA) (01°27'25 "S, 48°26'36" W) in Belém, Pará,
79 Brazil. Seeds were sown in polyethylene pots (15 x 25 x 0.05 cm) filled with low-fertility soil
80 (Ferralsol - pH, 4,2; organic matter, 18,80 g dm⁻³; P, 2 mg dm⁻³; K, 4 mg dm⁻³; Ca, 0,2 mmolc
81 dm⁻³; Ca+Mg, 0,3 mmolc dm⁻³; Al, 1,4 mmolc dm⁻³) and kept under greenhouse conditions.

82 2.2 Preparation of inoculum

83 *Pseudomonas fluorescens* (BRM-32111) and *Burkholderia pyrrocinia* (BRM-32113)
84 are currently stored and preserved in the *in vitro* collection of the Plant Protection Laboratory,
85 at the Federal Rural University of the Amazon. The bacterial isolates were cultured in solid 523
86 medium for 48h at 28°C. The bacterial suspension was prepared in water and adjusted to A540
87 = 0.2 (10⁸ CFU/mL).

88 2.2 Inoculation forms tested

89 • *Seed* (microbiolized seed): *U. brizantha* seeds were sterilized with 70% ETOH
90 and 2% NaClO, both for 1 minute, washed in sterile water for 1 minute, and placed on sterile
91 filter paper, for 1 hour. Before sowing, the seeds were steeped in the suspensions for for 24
92 hours, at 28°C and at constant agitation.

93 • *Seed and Soil*: microbiolized seed + soil drenched.

94 • *Soil* (soil drenched): 5 mL of suspension of each treatment, bacterial isolates (10⁸
95 CFU) drenched the trial soil at 14 days after seedling emergence (DASE).

96 2.3 Experiment I

97 The treatments consisted of two microorganisms, BRM-32111 and BRM-32111 with
 98 three inoculation forms and a control. The experimental design was completely randomized
 99 with five replications. At 21 DASE, seedlings were harvested and separated into shoot (leaf
 100 blades and culms) and roots. Plant material was oven dried (60° C) until constant mass. Total
 101 dry mass (TDM) was calculated by adding shoot dry mass (SDM) and root dry mass (RDM).
 102 The experiment was repeated three times with similar results.

103 2.4 Experiment II

104 The treatments consisted of fertilized and unfertilized non-inoculated *B. brizantha*
 105 plants, or fertilized plants, inoculated with BRM-32111, BRM-32113 and co-inoculated with
 106 BRM-32111 + BRM-32113 (MIX). The experimental design was completely randomized with
 107 eight treatments and five replications. Fertilized plants were fertilized with 5 mg dm⁻³ of N, 14
 108 mg dm⁻³ of P₂O₅, and 10 mg dm⁻³ of K₂O. Suspension of bacterial isolates (5 mL, 10⁸ CFU),
 109 water drenched the trial soil at 14 DASE. At 21 DASE, seedlings were harvested to determine
 110 biomass production. The experiment was repeated three times with similar results.

111 2.5 Experiment III: Growth promotion effects of microorganism on *B. brizantha*

112 The treatments consisted of non-inoculated *B. brizantha* fertilized- (positive) and
 113 unfertilized- controls (negative), inoculated with BRM-32111, BRM-32113 and co-inoculated
 114 with (MIX). The experimental design was completely randomized with five replications and
 115 five treatments. The all experiment was conducted in a greenhouse, with mean air temperature
 116 of 30 ± 2.5 °C and relative humidity of 74% ± 4 (mean ± s.d.), respectively.

117 Plant growth parameters were calculated according to Hunt (1990) and Barbero et al.,
 118 (2013). At 35 DASE, plants were harvested to determine biomass production. We calculated
 119 root/shoot dry mass ratio (RDM/SDM). Leaf area (LA) was determined over leaf disks of either
 120 0,42 cm² or 2,28 cm², dried at 60°C until constant mass. The biomass allocation pattern was

121 estimated as the leaf, culm and root mass ratios (respectively, the ratio between total leaf, culm,
122 and root dry mass per plant and total dry mass per plant).

123 Five evaluation periods (14, 17, 21, 28 and 35 DASE) were used to determine the
124 number of leaves (NL), height (H), culm length (CL) and chlorophyll content (SPAD index -
125 soil plant analysis development), estimated by a portable chlorophyll meter (SPAD-502. Konica
126 Minolta Sensing, INC. Japan). Relative growth rate (change in total mass per total dry mass of
127 plant per day, RGR) was calculated for harvests at 14 and 35 DASE. Morphogenetic and
128 structural parameters, calculated according to Gomide and Gomide (2000), were: leaf
129 appearance rate (ratio between the difference in the number of initial and final leaves the
130 number of evaluation interval days, LApR), leaf elongation rate (ratio between the difference
131 of the initial and final lengths of the expanded sheets and the number of days of the evaluation
132 interval, LER), number of leaves per plant (NL) and number of tillers per plant (NT).

133 Net photosynthesis (A), stomatal conductance (g_s) and transpiration (E) were measured
134 35 days after seedling emergence, on one young, fully expanded blade per plant, with an
135 infrared gas analyzer (IRGA) (LI-6400XT; LICOR, Lincoln, NE). Measurements were made
136 under CO_2 of $400 \mu\text{mol m}^{-1}$ and a constant photosynthetic active radiation of $1000 \mu\text{mol m}^{-2}\text{s}^{-1}$
137 (obtained by an artificial light source coupled to the IRGA chamber).

138 2.4 Biochemical assays

139 For determination of free ammonium, nitrate, amino acid, total soluble proteins, and
140 mineral analysis of nitrogen (N) we selected the treatment that promoted the greatest growth in
141 *B. brizantha* (co-inoculated with BRM-32111 + BRM-32113) and two non-inoculated controls.

142 For determination of the free ammonium, 50 mg of dry matter incubated with 5 mL of
143 sterile distilled water at 100°C for 30 min, and was centrifuged at 2.000 g for 5 min at 20°C and
144 the supernatant was removed. The quantification of the free ammonium was carried out at 625
145 nm in accordance with Weatherburn (1967), with $(\text{NH}_4)_2\text{SO}_4$ as standard.

For determination of nitrate, 100 mg of dry matter was incubated with 5 ml of sterile distilled water at 100°C for 30 min. The homogenized mixture was centrifuged at 3.000 g for 15 min at 25°C, and the supernatant was removed. The quantification of the nitrate was carried out at 410 nm in accordance to Cataldo et al., (1975), with KNO₃ as standard.

The amino acid was determined using the 50 µl ethanolic extract, 50 µl Na-Citrate (1M + 0,2% Ascorbic Acid (100mL NaCitrate + 0,2 g Asc. Acid)) and 100 µl Ninhydrin solution (1%). The mixture was incubated at 95°C for 20 min. and centrifuged at 12000g for 10s. Absorbance was measured at 570 nm. The calibration curve was made using Leucine (1mM) (Gibon et al., 2004). For total soluble proteins, each pellet was vigorously shaken in 1 mL absolute ethanol, incubated at 80 °C for 20 min, and centrifuged at 12000g for 5 min., at 4 °C. The supernatant was discarded and the pellet was shaken with 1 mL 0.2 M KOH. After heating for 60 min at 90 °C, samples were cooled and centrifuged at 12,000g, for 5 min, at 4 °C. Quantification of the total soluble proteins was carried out at 595 nm in accordance with Bradford (1976), with albumin bovine as standard. Shoot mineral analysis of nitrogen (N) was determined by inductively coupled plasma optical emission spectrometry (ICPOES).

2.5 Statistical analysis

All Data were subjected to analysis of variance and variables with significant F values were compared by Duncan test ($P < 0.05$). The LN, H, SL and SPAD were analyzed by ANOVA in a factorial arrangement (evaluation period x treatment). *Post hoc* contrasts were calculated for assessing differences between controls and inoculated plants for LN, H, SL, SPAD, MST, LA, *A*, *gs* and *E*. Parametric correlation analysis was calculated between SPAD, LA, H, NT, N or *A* versus MST. The statistical package STATISTICA for Windows release 7 (StatSoft, Inc., Tulsa, USA) was used for all computations of the data.

3. Results

3.1 Inoculation forms tested

171 No increase in biomass production could be observed when microorganisms were
 172 inoculated solely in the seeds or seed + soil drench (Table 1). When inoculation was performed
 173 by soil drench *P. fluorescens* (BRM-32111) and *B. pyrrocinia* (BRM-32113) increased biomass
 174 production ($F_{1,36} = 1989,23$; $P < 0.01$) by 242% and 112%, respectively (Table 1).

175 3.2 Effects of fertilizers and PGPR on *B. brizantha*

176 No increase in biomass production could be observed when microorganisms were
 177 inoculated on fertilized plants (Table 2). However, inoculation of unfertilized plants with BRM-
 178 32113, BRM-32111 or Mix (BRM-32111 + BRM-32113) increased biomass production by
 179 95%, 227% and 327%, respectively, relative to non-inoculated unfertilized-control plants ($F_{1,32}$
 180 $= 3123,7$; $P < 0.01$) (Table 2).

181 3.3 Growth promotion effects of microorganism on *B. brizantha*

182 The total number of leaves ($F_{1,75} = 537.6$; $P < 0.01$), plant height ($F_{1,75} = 2387.2$; $P <$
 183 0.01), culm length ($F_{1,75} = 241.5$; $P < 0.01$) and Spad index ($F_{1,75} = 963.9$; $P < 0.01$) were higher
 184 in inoculated plants (Fig. 1 and 2). The beneficial effects of rhizobacteria on *B. brizantha*
 185 development could already be observed three days after inoculation (17 DASE) (Fig. 2).

186 Leaf area was increased ($F_{1,20} = 186.9$, $P < 0.01$) by inoculation. This increment was
 187 above 700% relative to unfertilized-control plants and 108% relative to fertilized-control plants
 188 (Table 3). The RDM/ADM ratio was higher in co-inoculated plants (Table 3). The RGR ranged
 189 from 0.1 to 0.17, being higher in co-inoculated plants (Table 3).

190 The rhizobacteria increased NT and the mean EF (Table 3). The L/C ratio was higher in
 191 plants co-inoculated and unfertilized-control plants (Table 3). LApR and LER from inoculated
 192 plants were higher than those of the control plants, either fertilized, or unfertilized (Table 3).
 193 The photosynthetic rate ($F_{1,20} = 364.8$, $P < 0.01$), stomatal conductance ($F_{1,20} = 84.24$; $P < 0.01$)
 194 and transpiration ($F_{1,20} = 72.70$, $P < 0.01$) were higher in inoculated plants (Fig. 3).

195 Biomass production was higher in inoculated plants ($F_{1,20} = 2289.5$, $P < 0.01$), with an
 196 increment of shoot dry mass of more than 930%, relative to unfertilized-control plants, and over
 197 334% relative to fertilized-control plants (Fig. 4a). For root dry mass production, this increment
 198 was above 770% and 262% relative to unfertilized- and fertilized-control plants, respectively
 199 (Fig. 4b).

200 When the rhizobacteria were inoculated individually, the increment in total biomass
 201 production was over 870% and 300%, respectively, relative to unfertilized- and fertilized-
 202 control plants. The maximum gain in total biomass production was achieved by co-inoculation
 203 (MIX), which accounted for an increment of over 1300%, relative to unfertilized-control plants
 204 and nearly 500%, when compared to fertilized-control plants (Fig. 4c). Biomass allocation data
 205 revealed that, except for fertilized-control plants, in all treatments there was a preferential
 206 allocation to leaves (Fig. 4d). The increment in total biomass production correlated positively
 207 to the SPAD index, LA, H, NT and *A* (Table 4).

208 3.3 Biochemical effects of microorganism on *B. brizantha*

209 Relative to unfertilized- and fertilized-control plants, co-inoculated plants showed
 210 higher nitrate concentration in leaves (130% and 20%), roots (60% and 16%) and total biomass
 211 (100% and 18%); amino acid in the root (135%); protein concentration in the leaves (33% and
 212 12%), root (142% and 21) and total biomass (80% and 6%) (Fig. 5) and higher nitrogen
 213 concentration in leaves (30% and 11%), root (75% and 25%) and total biomass (27% and 19%)
 214 (Fig. 6). Higher concentrations of ammonium (leaf, root and total) and amino acids (leaves and
 215 total) were found in unfertilized-control plants (Fig. 5).

216 4. Discussion

217 *P. fluorescens* and *B. pyrrocinia* fostered the highest growth in *B. brizantha* cv. Piatã,
 218 when inoculated by soil drench, during seedling stage (Table 1). Failure to promote plant

219 growth, when these bacteria were inoculated in the seeds, may indicate that, during germination,
 220 *B. brizantha* might be able to recognize microbial compounds, synthesizing substances capable
 221 of inhibiting the beneficial effects of these rhizobacteria on plant growth promotion (Doornbos
 222 et al., 2012). A similar mechanism of plant immune stimulation probably was also activated,
 223 when plants were sequentially inoculated, both in the seed and by soil drench, inhibiting growth
 224 promotion, could be related to the allelopathic potential of *Urochloa* (Souza Filho et al., 2005;
 225 Kato-Noguchi et al., 2014). Under this condition, allelopathy can also affect the rhizosphere
 226 microbial community, and may be the cause of the observed lower *Bacillus* spp. colonization
 227 in *B. brizantha* rhizosphere (Araujo and Pedroso, 2013).

228 The amount allelopathic root exudates compounds may also vary during the plant's
 229 developmental stage (Doornbos et al., 2012). However, allelopathic compounds exudated by
 230 *Brachiaria* roots are known to have no inhibitory effects at low concentrations (Souza Filho et
 231 al., 2005). Thus, we can infer that the allelopathic compounds, detrimental to rhizobacteria,
 232 possibly exuded by *B. brizantha* roots in our study, might have decreased over time, because
 233 there was an increased growth of *B. brizantha* plants, inoculated by soil drench (Table 1).

234 Soil fertilization was antagonistic the bacterial activities on the growth of *B. brizantha*
 235 (Table 2). On the other hand, inoculation with rhizobacteria increased growth in unfertilized
 236 plants (i.e., exposed to nutrient limitation) (Table 2). It could be inferred that roots of
 237 unfertilized plants modified rhizodeposition patterns, by secreting specific compounds,
 238 resulting in an increased microbial biomass and activity around the roots (Doornbos et al., 2012;
 239 Carvalhais et al., 2013). In maize plants, the nutritional status affects the root colonizing
 240 bacterium, stimulating the repression of genes associated with protein synthesis, changing the
 241 composition of root exudates, and influencing the physiology of associative bacteria
 242 (Carvalhais et al., 2013). In temperate grasslands, Keuter et al., (2014) observed that
 243 fertilization decreases non-symbiotic biological N fixation, through the inhibition of

244 nitrogenase. The higher N-fertilizer doses also reduce of the beneficial bacterial effects on the
 245 growth in wheat (Saubidet et al., 2002; Silveira et al., 2016) and *Sorghum bicolor* plants (Santos
 246 et al., 2017).

247 Our results show the potential of *P. fluorescens* and *B. pyrrocinia* for increasing plant
 248 growth in *B. brizantha* cv. Piatã (Fig. 1). It seems that *P. fluorescens* and *B. pyrrocinia* probably
 249 acted synergistically in co-inoculated plants, as plant growth was higher when they were
 250 inoculated individually (Fig. 4). Increases in biomass production in *B. brizantha* were of over
 251 20% and 14%, after seed inoculation with *Bacillus* (Araujo et al., 2012) and *Azospirillum*
 252 *brasilense* (Hungria et al., 2016), and of over 100%, after root inoculation with endophytic
 253 bacteria (Kelemu et al., 2011).

254 Tillering, root development and a high root /shoot dry mass ratio are important features
 255 for an efficient pasture establishment. In the present study, these attributes increased in co-
 256 inoculated plants as result of an increased nitrogen concentration in roots. Greater tillering and
 257 root biomass were also reported for *Panicum virgatum* inoculated with *Burkholderia*
 258 *phytofirmans* (Kim et al., 2012). Inoculated plants developed a higher leaf length, area, and
 259 number, probably increasing their light capture ability. In addition, net photosynthesis,
 260 evaluated on an area basis, was enhanced by inoculation. These improved responses might have
 261 contributed to the increased relative growth rate and biomass production measured on those
 262 plants. The positive effect of plant growth-promotion rhizobacteria on net photosynthetic is also
 263 reported in rice (Nascente et al., 2016).

264 The relatively lower net photosynthetic rates of the co-inoculated plants, relative to
 265 plants individually inoculated, could be attributed to a likely more advanced physiological stage
 266 of these fast-growing, co-inoculated plants. In this regard, Wang et al., (2015) report a faster
 267 decline, with plant age, in the rates of photosynthesis, transpiration and stomatal conductance
 268 in *Panicum virgatum* inoculated with *Burkholderia phytofirman*. That is, *B. phytofirman*

269 accelerated development and maturation in *Panicum virgatum* seedlings, as well as induced
 270 earlier senescence and flowering in adult plants. According to Larcher (2006), gas exchange
 271 ability changes during plant development, tending to correlate negatively with the physiological
 272 stage. As the co-inoculated plants showed a higher number of tillers and relative growth rate,
 273 we could assume these plants were in a more advanced physiological stage.

274 In co-inoculated plats, the amino acids had rapid conversion into proteins, increasing
 275 nitrogen concentration and Spad index (chlorophyll content). This increased the development
 276 of photosynthetic organs, enhancing leaf length, leaf appearance rate, leaf area and biomass
 277 allocation to the leaves. These are desirable characteristics for forage grasses, since leaf blades
 278 are the preferred nutrient source for ruminants, for their higher protein content and digestibility.
 279 The inoculation with *Azospirillum brasilense* also promoted greater nitrogen uptake and biomass
 280 production in *B. brizantha* (Hungria et al., 2016) and in wheat plants (Saubidet et al., 2002;
 281 Silveira et al., 2016). Higher Spad index, nitrogen content and biomass production were also
 282 reported in *Brachiaria* with bacterial endophytes, under low nutrient conditions (Kelemu et al.,
 283 2011).

284 The increase in nitrate and nitrogen concentrations observed in the tissues of the
 285 inoculated plants is probably a response of organic matter mineralization by the rhizobacteria
 286 (Pii et al., 2015), followed by nitrification. It is possible that the rhizobacteria alters the nitrate
 287 fluxes at the root plasma membrane (Saubidet et al., 2002; Pii et al., 2015), decreases the nitrate
 288 concentration at the root cell surface (rhizosphere), stimulating root development and increasing
 289 nitrate uptake capacity (Mantelin and Touraine, 2003).

290 In forage grass, nitrate fertilization increases the protein contents and biomass
 291 production, but in excess can be toxic to cattle (0.35 to 0.45 dag/kg) (Corrêa et al., 2007; Costa
 292 et al., 2009). In our study, levels of nitrate in the leaves of co-inoculated plants did not reach
 293 toxic levels. In addition, the increase in nitrate concentration in *B. brizantha*, could improve its

294 resistance to spittlebug attacks, as a higher nitrate concentration in the xylem is known to impair
 295 spittlebug nymphal development (Thompson, 2004).

296 Our results showed that inoculation of rhizobacteria by soil drench, at seedling stage,
 297 enhanced beneficial morphological and physiological characteristics, and revealed a direct
 298 positive effect of plant growth-promoting rhizobacteria on biomass production of *B. brizantha*
 299 cv. Piatã, cultivated on a low-fertility soil. Because, nitrogen concentration and Spad index
 300 (chlorophyll content) was highly and positively correlated to total biomass production (Table
 301 3). In addition, this might have contributed to higher net photosynthesis found in inoculated
 302 plants (Table 3). The higher biomass production of inoculated plants also related to greater leaf
 303 area and the number of tillers (Table 3), which, in turn, we could infer, was possibly favored
 304 by an enhanced auxin biosynthesis in inoculated plants. This, relationship will be investigated
 305 in future studies on changes in the plant hormones pathway of inoculated *B. brizantha* plants.

306 Furthermore, based on our results, it could be stated that the method of inoculation is
 307 key in determining the potential of microorganism for promoting plant growth. This is because
 308 inoculation by soil drench, at seedling stage, was more effective than inoculation only in the
 309 seed or both in the seed and by soil drench. Also, the fertilizer may have suppressed the
 310 beneficial bacterial effects on the growth of *B. brizantha*. Our results attested that *P. fluorescens*
 311 and *B. pyrrocinia* could promote growth in *B. brizantha* cv. Piatã, especially when co-
 312 inoculated. Our results point out to a potentially valuable source of practical information in the
 313 search of an eco-friendlier approach to increase pasture productivity.

314 Acknowledgments

315 We thank FAPESPA (Fundação de Amparo à Pesquisa do Estado do Pará) for financial
 316 support. We also thank Francisco Janyelo Palacios Martinez, from Grupo Gasparim, for kindly
 317 supplying the *Brachiaria brizantha* cv. Piatã seeds, and Ana Carolina Sonsim de Oliveira Bueno
 318 and Marcela Cristiane Ferreira Rêgo for technical laboratory support.

319 References

- 320 Araujo, F.F., Guaberto, L.M., Silva, I.F., 2012. Bioprospecção de rizobactérias promotoras de
321 crescimento em *Brachiaria brizantha*. R. Bras. Zootec. 41, 521-527.
- 322 Araujo, F.F., Pedroso, R.A.B., 2013. Interação de *Bacillus* sp. com a rizosfera de três espécies
323 de plantas forrageiras. Biosci. J. 29, 152-158.
- 324 Barbero, L.M., Prado, T.F., Basso, K.C., Lima, L.A., Motta, K.M, Krüger, B.C., Martins Neto,
325 L.R., Silva, G.A.S., 2013. Análise de crescimento em plantas forrageiras aplicada ao
326 manejo de pastagens. Vet. Not. 19, 71-85.
- 327 Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram
328 quantities of protein utilizing the principle of protein-dye binding. Anal Biochem. 72, 248-
329 254.
- 330 Carvalhais, L.C, Dennis, P.G, Fan, B., Fedoseyenko, D., Kierul, K., Becker A., Wiren, N.,
331 Borriss, R., 2013. Linking Plant Nutritional Status to Plant-Microbe Interactions. PLoS
332 ONE 8, (7), e68555.
- 333 Cataldo, D.A., Haroon, S.L.E., Yougs, V.L., 1975. Rapid colorimetric determination of nitrate
334 in plant tissue by nitration of salicylic acid. Commun Soil Scie Plant Anal. 6, 71-80.
- 335 Corrêa, L.A., Cantarella, H., Primavesi, A.C., Primavesi, O., Freitas, A.R., Silva A.G., 2007.
336 Efeito de fontes e doses de nitrogênio na produção e qualidade da forragem de capim-
337 coastcross. R. Bras. Zootec., 36, 763-772.
- 338 Costa, K.A.P., Oliveira, I.P., Faquin, V., Silva, G.P., Severiano, E.C., 2009. Produção de massa
339 seca e nutrição nitrogenada de cultivares de *Brachiaria brizantha* (a. Rich) stapf sob doses
340 de nitrogênio. Ciênc. agrotec., 33, 1578-1585.
- 341 Dias-Filho, M.B., 2011. Degradação de pastagens: processos, causas e estratégias de
342 recuperação. 4. Ed., rev. atual. e ampl. Belém, PA.
- 343 Doornbos, R.F, Loon L.C.V., Bakker, P.A.H.M., 2012. Impact of root exudates and plant
344 defense signaling on bacterial communities in the rhizosphere. A review. Agron. Sustain.
345 Dev. 32, 227-243.
- 346 Gibon, Y., Blaessing, O.E., Hannemann, J., Carillo, P., Hohne, M., Hendriks, J.H., Palacios, N.,
347 Cross, J., Selbig, J., Stitt, M., 2004. A Robot-based platform to measure multiple enzyme
348 activities in Arabidopsis using a set of cycling assays: comparison of changes of enzyme
349 activities and transcript levels during diurnal cycles and in prolonged darkness. Plant Cell
350 16, 3304-3325.
- 351 Gomide, C.A.M., Gomide, J.A., 2000. Morfogênese de cultivares de *Panicum maximum* Jacq.
352 R. Bras. Zootec. 29, 341-348.
- 353 Hungria, M., Nogueira, M.A., Araujo, R.S., 2016. Inoculation of *Brachiaria* spp. with the plant
354 growth-promoting bacterium *Azospirillum brasilense*: An environment-friendly
355 component in the reclamation of degraded pastures in the tropics. Agr. Ecosyst. Environ.
356 221, 125-131.
- 357 Hunt, R., 1990. Basic growth analysis for beginners, London.
- 358 Kato-Noguchi, H., Kobayashi, A., Ohno, O., Kimura, F., Fujii, Y., Suenaga, K., 2014.
359 Phytotoxic substances with allelopathic activity may be central to the strong invasive
360 potential of *Brachiaria brizantha*. J. Plant. Physio. 171, 525-530.
- 361 Kelemu, S., Fory, P., Zuleta, C., Ricaurte, J., Rao, I., Lascano, C., 2011. Detecting bacterial
362 endophytes in tropical grasses of the *Brachiaria* genus and determining their role in
363 improving plant growth. African J. Biotech. 10, 965-976.
- 364 Keuter, A., Veldkamp, E., Corre, M.D. 2014 Asymbiotic biological nitrogen fixation in a
365 temperate grassland as affected by management practices. Soil Biol. Biochem. 70, 38-46.

- Kim, S., Lowman, S., Hou, G., Nowak, J., Flinn, B., Mei, C., 2012. Growth promotion and colonization of switchgrass (*Panicum virgatum*) cv. Alamo by bacterial endophyte *Burkholderia phytofirmans* strain PsJN. *Biotech. Biof.* 5, 1-10.
- Larcher, W., 2006. *Ecofisiologia vegetal*. São Carlos, RIMA Artes e Textos.
- Mantelin, S., Touraine, B. 2003. Plant growth-promoting bacteria and nitrate availability: impacts on root development and nitrate uptake. *J. Exp. Bot.* 1-8.
- Marques, A.C.R., Oliveira, L.B., Nicoloso, F.T., Jacques, R.J.S., Giacomini, S.J., Quadros, F.L.F., 2017. Biological nitrogen fixation in C4 grasses of different growth strategies of South America natural grasslands. *Appl. Soil Ecol.* 113, 54 – 62.
- Nascente, A.S., Filippi, M.C.C., Lanna, A.C., Souza, A.C.A., Lobo, V.L.S., Silva, G.B., 2016. Biomass, gas exchange, and nutrient contents in upland rice plants affected by application forms of microorganism growth promoters. *Environ. Sci. Pollut. Res.* 24, 2956.
- Pii, Y., Mimmo T., Tomasi N., Terzano R., Cesco S., Crecchio C. (2015) Microbial interactions in the rhizosphere: beneficial influences of plant growth-promoting rhizobacteria on nutrient acquisition process. A review. *Biol Fertil Soils.* 51, 403.
- Rêgo, M.C.F., Borges, F.I., Filippi, M.C.C., Gonçalves, L.A., Silva, G.B., 2014. Morphoanatomical and Biochemical Changes in the Roots of Rice Plants Induced by Plant Growth-Promoting Microorganisms. *J. Botany.*
- Santos, C.L.R., Alves, G.C., Macedo, A.V.M., Giori, F.G., Pereira, W., Urquiaga, S., Reis, V.M., 2017. Contribution of a mixed inoculant containing strains of *Burkholderia* spp. and *Herbaspirillum* ssp. to the growth of three sorghum genotypes under increased nitrogen fertilization levels. *Appl. Soil Ecol.* 113, 96–106.
- Saubidet, M.I., Fatta, N., Barneix, A.J., 2002. The effect of inoculation with *Azospirillum brasilense* on growth and nitrogen utilization by wheat plants. *Plant and Soil* 245, 215–222.
- Silveira, A.P.D., Sala, V.M.R., Cardoso, E.J.B.N., Labanca, E.G., Cipriano, M.A.P., 2016. Nitrogen metabolism and growth of wheat plant under diazotrophic endophytic bacteria inoculation. *Appl. Soil Ecol.* 107, 313–319.
- Souza Filho, A.P.S., Pereira, A.A.G., Bayma, J.C., 2005. Aleloquímico produzido pela gramínea forrageira *Brachiaria humidicola*. *Planta Daninha.* 23, 25-32.
- Strassburg, B.B.N., Latawiec, A.E., Barioni, L.G., Nobre, C.A., Silva, V.P., Valentim, J.F., Vianna, M., Assad, E.D., 2014. When enough should be enough: improving the use of current agricultural lands could meet production demands and spare natural habitats in Brazil. *Global Environ. Change.* 28, 84-97.
- Thompson, V., 2004. Associative nitrogen fixation, C4 photosynthesis, and the evolution of spittlebugs (Hemiptera: Cercopidae) as major pests of neotropical sugarcane and forage grasses. *Bull. Entomol. Res.* 94, 189–200.
- Timmusk, S., Behers, L., Muthoni, J., Muraya, A., Aronsson, A.C., 2017. Perspectives and challenges of microbial application for crop improvement. *Frontiers in Plant Science.* 8, 49.
- Valle, C.B., Barrios, S.C.L., Jank, L., Santos, M.F., 2014. Melhoramento de plantas forrageiras para uma pecuária de baixa emissão de carbono. In: Pedreira, B.C., Pereira, D.H., Pina, D.S., Carnevali, R.A., Lopes, L.B. *Intensificação da produção animal em pastagens*. Brasília, DF: Embrapa, pp 109-139.
- Wang, B.; Seiler, J.R.; Mei, C., 2015. *Burkholderia phytofirmans* strain PsJN advanced development and altered leaf level physiology of switchgrass. *Biomass and Bioenergy* 83, p. 493-500.
- Weatherburn, M.W., 1967. Phenol hypochlorite reaction for determination of ammonia. *Analytical Chemistry.* 39, 971-974.

415 Tables

Table 1 Inoculation method of plant growth-promoting rhizobacteria on the biomass production of *Brachiaria brizantha*. Shoot dry mass (SDM), root dry mass (RDM), total dry mass (TDM).

Inoculation	Isolates	Biomass (mg)		
		SDM	RDM	TDM
Seed	<i>Pseudomonas</i> spp.	54 ± 1.03 c	29.6 ± 0.89 c	83.6 ± 1.71 c
	<i>Burkholderia</i> spp.	50.9 ± 1.77 c	28.2 ± 0.80 cd	79.1 ± 2.46 c
Seed	<i>Pseudomonas</i> spp.	55.7 ± 1.01 c	26.8 ± 0.97 d	82.5 ± 1.08 c
+ Soil	<i>Burkholderia</i> spp.	53.6 ± 0.51 c	29.8 ± 0.93 c	83.4 ± 1.36 c
	<i>Pseudomonas</i> spp.	202.2 ± 0.81 a	89.2 ± 1.25 a	291.4 ± 1.89 a
Soil	<i>Burkholderia</i> spp.	123.2 ± 1.21 b	57.6 ± 1.44 b	180.8 ± 1.93 b
	Control	54.6 ± 1.67 c	30.8 ± 0.77 c	85.4 ± 1.14 c

* Significant at the 0.05 probability level. Data are means ± SE. Means followed by different letters in each column are significantly different ($P < 0.05$, Duncan Test).

Table 2 Effects of fertilization and plant growth-promoting rhizobacteria on the biomass production of *Brachiaria brizantha*. Shoot dry mass (SDM), root dry mass (RDM), total dry mass (TDM).

Treatments		Biomass (mg)		
		SDM	RDM	TDM
Fertilized	<i>Pseudomonas</i> spp.	104.6 ± 0.68 d	47.8 ± 0.37 d	150.8 ± 0.91 d
	<i>Burkholderia</i> spp.	103.1 ± 0.71 d	48 ± 0.32 d	150.8 ± 0.58 d
	Mix	102.8 ± 0.73 d	47.8 ± 0.42 d	150.4 ± 1.23 d
	Control	102.6 ± 0.76 d	47.6 ± 0.75 d	152.2 ± 1.39 d
Unfertilized	<i>Pseudomonas</i> spp.	203.8 ± 1.35 b	87 ± 0.83 b	290.8 ± 2.18 b
	<i>Burkholderia</i> spp.	116.4 ± 1.03 c	56.6 ± 0.87 c	173 ± 1.82 c
	Mix	282.4 ± 2.11 a	96.6 ± 0.93 a	379 ± 2.87 a
	Control	58.4 ± 0.81 e	30.4 ± 0.67 e	88.8 ± 1.24 e

* Significant at the 0.05 probability level. Data are means ± SE ($n = 5$). Means followed by different letters in each column are significantly different ($P < 0.05$, Duncan Test).

Table 3 Leaf area (LA cm²), root dry mass (RDM - g day⁻¹) /shoot dry mass (SDM) ratio, relative growth rate (RGR), number of tillers (NT), expanded leaf length (EF - cm), leaf appearance rate (LApR - L⁻¹ day⁻¹) and leaf elongation rate (LER - cm day⁻¹) of *Brachiaria brizantha* with growth-promoting rhizobacteria.

Treatments	LA	RDM /SDM	RGR	NT	EF	LApR	LER
C -	19.02 ± 1.96 e	0.65 ± 0.02 b	0.01 ± 0.0004 e	0 ± 0 d	22.74 ± 0.59 d	0.19 ± 0.02 e	0.90 ± 0.08 e
C +	73.70 ± 13.27 d	0.64 ± 0.02 b	0.03 ± 0.0003 d	1 ± 0 c	37.28 ± 0.56 c	0.27 ± 0.02 d	1.40 ± 0.10 d
B	187.20 ± 9.57 b	0.54 ± 0.01 c	0.12 ± 0.0042 c	2 ± 0 b	45.66 ± 0.50 ab	0.49 ± 0.05 c	1.84 ± 0.10 c
P	153.51 ± 7.74 c	0.62 ± 0.02 b	0.13 ± 0.0017 b	2 ± 0 b	43.54 ± 0.72 b	0.66 ± 0.04 b	2.53 ± 0.19 b
MIX	334.05 ± 25.58 a	0.68 ± 0.02 a	0.17 ± 0.0036 a	4 ± 0 a	46.72 ± 0.53 a	0.71 ± 0.05 a	2.79 ± 0.24 a

*Significant at the 0.05 probability level. Data are means ± SE (n = 5). Means followed by different letters in each column are significantly different (P < 0.05, Duncan Test).

† C - = unfertilized-control; C + = fertilized-control; B = *Burkholderia pyrrocinia*; P = *Pseudomonas fluorescens*; MIX = B + P.


Table 4 Correlation coefficient (r) of the correlations between SPAD, leaf area (LA), plant height (H), number of tillers (NT), photosynthesis (A) and nitrogen concentration (N) versus total dry mass (TDM) of *Brachiaria brizantha*.

Parameters	r
SPAD	0.84*
LA	0.95*
H	0.90*
NT	0.96*
A	0.60*
N	0.89*

*P < 0.05.

447

Figures



448

449

450 **Fig. 1** Shoot (a-e) and root (f-j) of *Brachiaria brizantha*, 21 days after inoculation (35 days after
 451 seedling emergence). Unfertilized-control (a, f), fertilized-control (b, g), inoculated with
 452 *Pseudomonas fluorescens* (BRM-32111) (c, h), *Burkholderia pyrrocinia* (BRM-32113) (d, i)
 453 and co-inoculated with BRM-32111 + BRM-32113 (e, j).

454

455

456

457

458

459




Fig. 2 Number of leaves per plant (a), plant height (H) (b), culm length (c) and SPAD index (d) of *Brachiaria brizantha* inoculated with growth-promoting rhizobacteria. Values are means $\pm$ SE ($n = 5$). Days = days after sowing. C - = unfertilized-control; C + = fertilized-control; B = *Burkholderia pyrrocinia*; P = *Pseudomonas fluorescens*; MIX = B + P.





Fig. 3 *Brachiaria brizantha* responses to growth-promoting rhizobacteria. (a) Net photosynthesis (A), (b) stomatal conductance (gs) and (c) transpiration (E). At 21 days after inoculation (35 days after seedling emergence). Columns with different letters are significantly different among treatments (P < 0.05, Duncan Test). C - = unfertilized-control; C + = fertilized-control; B = *Burkholderia pyrrocinia*; P = *Pseudomonas fluorescens*; MIX = B + P.



490

491 **Fig. 4** *Brachiaria brizantha* responses to growth-promoting rhizobacteria. (a) Shoot dry mass
 492 production (SDM), (b) root dry mass production (RDM), (c) total dry mass production (TDM)
 493 and, (d) biomass allocation pattern. 21 days after inoculation (35 days after seedling
 494 emergence). Columns with different letters are significantly different among treatments ($P <$
 495 0.05, Duncan Test). Different upper-case letters within columns indicate significant differences
 496 among plant organs ($P <$ 0.05, Duncan Test). C - = unfertilized-control; C + = fertilized-control;
 497 B = *Burkholderia pyrrocinia*; P = *Pseudomonas fluorescens*; MIX = B + P.


498

499

500

501


502



503

504 **Fig. 5** *Brachiaria brizantha* responses to growth-promoting rhizobacteria. (a) Free ammonium,
 505 (b) nitrate, (c) amino acid and, (d) total soluble proteins. 21 days after inoculation (35 days after
 506 seedling emergence). Columns with different letters are significantly different among
 507 treatments ($P < 0.05$, Duncan Test). C - = unfertilized-control; C + = fertilized-control; Mix=
 508 *Burkholderia pyrrocinia* + *Pseudomonas fluorescens*.

509



510

511 **Fig. 6** Nitrogen concentration of *Brachiaria brizantha* inoculated with growth-promoting
 512 rhizobacteria. 21 days after inoculation (35 days after seedling emergence). Columns with
 513 different letters are significantly different among treatments ($P < 0.05$, Duncan Test). C - =
 514 unfertilized-control; C + = fertilized-control; Mix= *Burkholderia pyrrocinia* + *Pseudomonas*
 515 *fluorescens*.

516

517 **Valor nutritivo, produção de biomassa e parâmetros anatômicos de *Brachiaria brizantha***
 518 **com rizobactérias promotoras de crescimento (PGPR)²**

519 Monyck Jeane dos Santos Lopes, Moacyr Bernardino Dias Filho, Thomaz Henrique dos Reis
 520 Castro, Edilson Freitas da Silva, Gisele Barata da Silva

521 **Resumo**

522 O uso de microrganismos promotores de crescimento é uma alternativa promissora e
 523 sustentável para aumentar a produtividade de pastagens. O objetivo foi avaliar o efeito da co-
 524 inoculação de *Pseudomonas fluorescens* (BRM- 32111) e *Burkholderia pyrrocinia* (BRM-
 525 32113) na anatomia, bioquímica, qualidade nutricional e produção de biomassa de *B. brizantha*.
 526 O delineamento experimental foi inteiramente casualizado com três tratamentos: plantas não
 527 inoculadas sem adubação (C-), com adubação (C +) e plantas coinoculadas com *P. fluorescens*
 528 e *B. pyrrocinia*. Foram avaliados parâmetros anatômicos, fisiológicos e de qualidade
 529 nutricional. A inoculação com rizobactérias modificou a anatomia foliar, do colmo e da raiz;
 530 incrementou o teor de clorofila (122%), taxas fotossintéticas (14%), eficiência do uso da água
 531 (22%), carboidratos totais (29%), amido (80%) e proteína bruta (37%); também aumentou os
 532 teores de N (30%), P (20%), Mg (20%) e Fe (180%), promovendo uma maior qualidade e
 533 produção de biomassa (400%) em *B. brizantha*. Por tanto, essas rizobactérias poderiam ser
 534 usadas como biofertilizantes para *B. brizantha*, contribuindo para a obtenção de forragem de
 535 maior produtividade e valor nutritivo.

536 **Palavras-chave:** biofertilizante, nitrogênio, fósforo, *Pseudomonas fluorescens*,
 537 *Burkholderia pyrrocinia*

² Este capítulo segue as normas de formatação da **Grass and Forage Science**.

538 **INTRODUÇÃO**

539 Na América Latina, o interesse crescente pelo sequestro de carbono e redução de gases
540 do efeito estufa, requer uma melhor gestão nos sistemas de pastagem (Pagano *et al.* 2017). As
541 pastagens tropicais são formadas principalmente por *Brachiaria* (Syn. *Urochloa*), uma
542 gramínea forrageira perene, com elevado crescimento, tolerância ao pastejo e alta
543 disponibilidade de forragem (Reis *et al.*, 2013; Pontes *et al.*, 2016; Pagano *et al.*, 2017). Dentre
544 as cultivares de *B. brizantha* disponíveis no mercado, destaca-se a BRS Piatã, por apresentar
545 menor alongamento da haste e sazonalidade na produção, maior acúmulo de forragem durante
546 o período seco e desempenho animal, e ser mais tolerante à cigarrinha-das-pastagens em relação
547 as outras cultivares dessa espécie (Reis *et al.*, 2013).

548 O uso de microrganismos promotores de crescimento seria uma solução sustentável e
549 promissora para melhorar a tolerância ao estresse biótico e abiótico, nutrição e produção de
550 pastagens (Pagano *et al.*, 2017; Timmusk *et al.*, 2017; Duchene *et al.*, 2017). Rizobactérias
551 promotoras de crescimento (PGPR) contribuem para o desenvolvimento das plantas através de
552 mecanismos diretos, estimulando a produção de fitormônios e atuando como biofertilizantes; e
553 de mecanismos indiretos, aumentando a resistência das plantas a estresses bióticos e abióticos.
554 As PGPR atuam como biofertilizantes por aumentar a disponibilidade de nutrientes no solo, a
555 biomassa, área radicular e a capacidade de absorção de nutrientes da planta, podendo afetar os
556 mecanismos bioquímicos e o estado nutricional, aumentando o crescimento e o rendimento das
557 plantas (Richardson *et al.*, 2009; Calvo *et al.*, 2014; Pii *et al.*, 2015; Timmusk *et al.*, 2017).

558 Estudos prévios demonstraram que a co-inoculação de *Pseudomonas fluorescens* e
559 *Burkholderia pyrrocinia* promovem o crescimento em *Brachiaria brizantha* (pesquisa
560 exploratória conduzida pelo primeiro autor). Nossa hipótese é que essas rizobactérias são
561 capazes de elevar tanto a quantidade, como a qualidade de forragem de *B. brizantha*, podendo

ser usadas como biofertilizantes. Por tanto, nosso objetivo foi avaliar o efeito da co-inoculação de *B. pyrrocinia* e *P. fluorescens* na anatomia, bioquímica, qualidade nutricional e produção de biomassa em *B. brizantha* cv. BRS Piatã.

MATERIAL E MÉTODOS

O experimento foi conduzido no Laboratório de Proteção de Plantas (LPP) e em casa de vegetação na Universidade Federal Rural da Amazônia (UFRA) (01°27'25"S, 48°26'36"W) em Belém, Pará, Brasil. O experimento foi inteiramente casualizado com três tratamentos: plantas controles não inoculadas sem fertilizante (C-), com fertilizante (C+) e coinoculadas com *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113), com dez repetições cada, cada repetição correspondente a uma planta. As rizobactérias foram repicadas em placas de Petri com meio de cultura 523 (Kado and Heskett, 1970) e incubadas durante 48 horas a 28°C. A suspensão bacteriana foi preparada com água estéril e ajustada a $540 \text{ nm} = 0.2$ (10^{-8} UFC/mL).

Foram semeadas dez sementes por vaso de polietileno (15 x 25 x 0.05 cm) em solo de baixa fertilidade (Ferralsol - pH, 4,2; organic matter, 18,80 g dm⁻³; P, 2 mg dm⁻³; K, 4 mg dm⁻³; Ca, 0,2 mmolc dm⁻³; Ca+Mg, 0,3 mmolc dm⁻³; Al, 1,4 mmolc dm⁻³). As plantas controles não inoculadas com fertilizante foram adubadas com 5 mg dm⁻³ de N, 14 mg dm⁻³ de P₂O₅ e 10 mg dm⁻³ de K₂O. A suspensão bacteriana (5 mL, 10⁸ CFU/mL) foi inoculada por rega do solo 14 dias após a emergência da planta (DAEP). O experimento foi conduzido em casa de vegetação, com radiação fotossinteticamente ativa de 900 $\mu\text{mol m}^{-2} \text{ s}^{-1}$, temperatura média do ar de 30 °C e umidade relativa de 74%.

A taxa de assimilação líquida de CO₂ (A) foi determinada 35 DAEP, com um analisador de gás infravermelho (IRGA) (LI-6400XT, LICOR, Lincoln, NE), com 400 $\mu\text{mol m}^{-1}$ de CO₂ e 1000 $\mu\text{mol m}^{-2} \text{ s}^{-1}$ de radiação fotossintética ativa (obtida por uma fonte de luz artificial acoplada

586 à câmara IRGA). A eficiência de uso da água (WUE) foi calculada pela razão entre a
587 fotossíntese e a transpiração. O teor de clorofila foi estimado com um medidor portátil (SPAD
588 - soil plant analysis development; SPAD-502. Konica Minolta Sensing, INC. Japão).

589 Aos 35 DAEP, as amostras de folhas, colmo e raízes foram coletadas e fixadas em
590 solução de FAA 50% (etanol: ácido acético glacial: formaldeído). O estudo anatômico foi
591 realizado no Laboratório de Anatomia Vegetal, Departamento de Botânica, Museu Emilio
592 Goeldi, MPEG. As amostras foram desidratadas em uma série de etanol graduada e emblocadas
593 em metacrilato de hidroxietilo (Leica®, Alemanha). Os blocos foram seccionados
594 transversalmente em um micrótomo rotativo Leica RM 2265. As secções foram coradas com
595 0,05% de azul de toluidina em tampão acetato, pH 4,3 (O'brien et al., 1964) e montadas em
596 resina sintética Entellan®.

597 As observações foram realizadas no microscópio Olympus BX61 (Japão) com câmera
598 digital acoplada conectada a um computador com software Motic 2.0. As medidas foram
599 realizadas usando o programa ImageJ para análise de imagem (Cappellari et al., 2015). Os
600 parâmetros anatômicos medidos foram: folhas - espessura da epiderme adaxial e abaxial, área
601 das células buliformes, do xilema, do floema, do esclerênquima, dos feixes vasculares, da
602 bainha do feixe e do mesófilo clorofiliano; colmo – espessura da epiderme, área e número de
603 feixes vasculares; e raiz - epiderme, exoderme, parênquima, córtex, endoderme, periciclo,
604 cilindro vascular, números de metaxilema e protoxilema e diâmetro do xilema. Para as
605 avaliações anatômicas, o delineamento experimental foi inteiramente casualizado, com dez
606 repetições por tratamento, cada repetição correspondente a uma planta, composto do valor
607 médio de três lâminas com seis secções cada.

608 Aos 35 DAEP, mensurou-se a altura (H) e as mudas foram coletadas, separando parte
609 aérea (folha e colmo) e raiz. O material vegetal coletado foi seco em estufa com temperatura de

610 60 °C até atingir massa constante. Para avaliar a produção total de biomassa, somou-se a massa
611 seca da parte aérea e raiz de cada planta. A área radicular foi estimada usando o programa de
612 imagem ImageJ (Tajima and Kato, 2011; Schneider *et al.* 2012). Os nutrientes N, C, P, K, Mg,
613 Ca, B, Fe, Zn e Cu foram determinados por espectrometria de emissão óptica com plasma
614 acoplado indutivamente (ICPOES). A fibra em detergente neutro e a fibra em detergente ácido
615 (base na matéria seca) foram determinadas utilizando o procedimento descrito por Van Soest *et*
616 *al.*, (1991). O teor de N foi analisado de acordo com o Kjeldahl. A proteína bruta foi calculada
617 multiplicando o teor de N (%) por 6.25. A extração dos pigmentos fotossintéticos foi em etanol
618 segundo Porra *et al.* (1989) e quantificado de acordo com Lichtenthaler (1987). Os carboidratos
619 solúveis totais e o amido foram determinados de acordo com Dubois *et al.* (1956).

620 Os dados foram submetidos a análise de variância e as variáveis com valores
621 significativos de F foram comparadas pelo teste de Duncan ($P < 0,05$). O software usado foi o
622 STATISTICA 7 (StatSoft, Inc., Tulsa, EUA)

623 **RESULTADOS**

624 **Parâmetros anatômicos**

625 O limbo foliar das plantas inoculadas com rizobactérias apresentou epiderme adaxial
626 com maior espessura (80%), e o aumento das áreas de células buliformes (30%), bainha dos
627 feixes (20%), feixes vasculares e mesófilo clorofiliano (10%) (Tabela 1; Figura 1 (a - c)). As
628 plantas adubadas apresentaram maior epiderme abaxial, área de esclerênquima e número de
629 feixes (Tabela 1; Figura 1 (a - c)).

630 O colmo das plantas com rizobactérias apresentou incremento em média de 900% na
631 quantidade de feixes vasculares em relação as plantas controles sem adubo e de 102%
632 comparado com as plantas controle adubadas (Tabela 1; Figura 1). Com floema voltado para

633 sua periferia e o xilema em oposição, também apresentou maior número de feixes. As plantas
634 com rizobactérias foram as únicas com anel esclerenquimático, apresentando 2-5 camadas de
635 células e medula oca (Figura 1f).

636 As rizobactérias modificaram a anatomia das raízes das plantas, incrementando em mais
637 de 50% a exoderme, periciclo, parênquima e cilindro vascular, de 30% o protoxilema e o
638 diâmetro das células do xilema e de 10% a endoderme em comparação as plantas controles
639 (Tabela 1; Figura 1 (g - i)).

640 **Qualidade nutritiva**

641 Nas folhas das plantas inoculadas os maiores incrementos foram nos teores de N (30%
642 e 11%) e Fe (180% e 150%) em comparação as plantas controle sem e com adubo (Tabela 2).
643 Também apresentaram um incremento médio de 20% do P, Na, e Mg em relação as plantas
644 controle; e de 23% em Zn e 9% em Cu, em comparação as plantas controle sem adubo. Nas
645 raízes das plantas inoculadas houve maior concentração de N, P, K, Na, Mg, Fe e Mg do que
646 as plantas controles (Tabela 2).

647 A quantidade de NDF não diferiu entre os tratamentos, e a de ADF foi maior nas plantas
648 controles com adubo (Tabela 2). O teor de proteína bruta (CP) foi maior nas plantas inoculadas,
649 com incremento em 37% em comparação as plantas controle sem adubo e 18% as plantas
650 adubadas (Tabela 2).

651 **Spad, trocas gasosas e parâmetros bioquímicos**

652 As plantas com rizobactérias apresentaram maiores concentrações de clorofila com
653 incremento médio de 143% em clorofila a, 95% em clorofila b e 122% no total (Figura 2 a). O
654 índice Spad aumentou em 37% em relação as plantas controles sem adubo e 11% em
655 comparação as plantas controles adubadas (Figura 2 b). As rizobactérias também

656 incrementaram em média de 14% a taxa fotossintética (*A*) e em 22% a eficiência momentânea
657 do uso da água (WUE) em relação as plantas controles (Figura 2 c, d).

658 O uso das rizobactérias incrementou a concentração de amido nas folhas (127% e
659 180%), nas raízes (46% e 73%) e o total (80% e 133%), em comparação com as plantas
660 controles sem e com adubo (Figura 3 a). Plantas inoculadas também apresentaram maior
661 concentração de carboidratos solúveis totais, com incremento de 45% nas folhas, 25% nas
662 raízes e 29% nos totais, em comparação as plantas controles (Figura 3 b).

663 Crescimento e produção de biomassa

664 As rizobactérias promoveram maior altura (mais de 100% e 30%), área radicular (131%
665 e 122%), produção de biomassa foliar (470%, e 450%), do colmo (480% e 79%), da raiz (330%
666 e 200%) e total (mais de 400% e 200%), em comparação as plantas controle sem e com adubo
667 (Figura 4).

668 DISCUSSÃO

669 A co-inoculação com *P. fluorescens* (BRM-32111) e *B. pyrrocinia* (BRM-32113)
670 modificou a anatomia, a bioquímica e o acúmulo de nutrientes, promovendo um maior
671 desenvolvimento em *B. brizantha* (Figura 5). Essas rizobactérias provavelmente aumentaram a
672 disponibilidade de nutrientes na rizosfera, através da fixação assimbiótica de nitrogênio;
673 solubilização de fósforo, por ácidos orgânicos e fosfatases; sequestro de ferro por produção de
674 sideróforos e liberação de micronutrientes de fontes insolúveis, além de estimular os sistemas
675 de transporte iônico na raiz (Vancheron *et al.*, 2013; Calvo *et al.*, 2014; Pii *et al.*, 2015).

676 A maior absorção de nutrientes pelas plantas inoculadas, também foi em resultado das
677 alterações que as rizobactérias causaram nas raízes, com o incremento da exoderme, reduzindo
678 o refluxo de íons para solução do solo, o que resultou em maior concentração de K e Mg nas

679 raízes; maior periciclo, resultante provavelmente do aumento de auxina, facilitando a difusão
680 de oxigênio da base para o ápice, e maior desenvolvimento de xilema, protoxilema e área
681 radicular, aumentando a absorção e fluxo de água e nutrientes (Richardson *et al.*, 2009;
682 Vacheron *et al.*, 2013).

683 No presente estudo, o aumento dos teores de nutrientes nas plantas inoculadas,
684 sobretudo o nitrogênio, como também a maior área da bainha do feixe vascular e do mesófilo
685 clorofiliano nas folhas, aumentou o teor dos pigmentos fotossintéticos (clorofila a, b e total).
686 Por sua vez, em decorrência do maior desenvolvimento do sistema vascular da folha, colmo e
687 raiz, aumentou o transporte de água e solutos, resultando em uma maior eficiência
688 fotossintética, incrementando a concentração de carboidratos totais e de amido, que tem alta
689 digestibilidade sendo fonte de energia aos animais. O uso de rizobactérias promotoras de
690 crescimento também aumentou a clorofila, taxas fotossintéticas e eficiência do uso da água em
691 *Phaseolus coccineus* (Stefan *et al.*, 2013); pigmentos fotossintéticos em *Mentha piperita*
692 (Cappellari *et al.*, 2015) e carboidratos e proteína em *Zea mays* L. (El-Ghany *et al.*, 2015),
693 *Arachis hypogaea* (Mathivanan *et al.*, 2017) e *Coriandrum sativum* L. (Warwate *et al.*, 2017).

694 É possível supor que os benefícios resultantes da inoculação também aumentariam a
695 tolerância de *B. brizantha* ao pastejo e a ambientes com limitação de recursos hídricos. Isso por
696 que as plantas inoculadas, além de apresentarem a raiz e o sistema vascular mais desenvolvido,
697 maiores taxas fotossintéticas e acúmulo de amido, também tiveram maior eficiência no uso de
698 água, e incremento em áreas de células buliformes, que são responsáveis por enrolar as folhas,
699 reduzindo a área de transpiração (Reis *et al.*, 2013). Além disso, as plantas inoculadas
700 apresentaram o colmo com anel esclerenquimático, o que as tornariam mais resistentes ao
701 acamamento (Reis *et al.*, 2013). A inoculação com rizobactérias também modificou a anatomia,
702 promovendo maior capacidade de adaptação, por aumentar o desenvolvimento vascular em

703 *Triticum aestivum* L. (El-Afry *et al.*, 2012), e a densidade de pelos radiculares em
704 *Handroanthus impetiginosus* (Larraburu and Llorente, 2015).

705 Em nosso estudo, a inoculação com rizobactérias incrementou os macro e
706 micronutrientes nas folhas (N, P, Na, Mg, Fe e Cu) e raízes (N, P, K, Na, Mg, Fe, Cu e Mn),
707 melhorando a translocação e mobilização de nutrientes, aumentando a qualidade nutricional de
708 *B. brizantha*. O maior teor de nitrogênio também resultou no incremento da proteína bruta (CP),
709 que é um indicador da qualidade da forragem, a qual deve ser superior a 7%, para não reduzir
710 o consumo voluntário e a digestibilidade pelo gado (Reis *et al.*, 2013; Ball *et al.*, 2015). Essa
711 maior nutrição nitrogenada nas plantas inoculadas, também foi verificada nas avaliações do
712 índice Spad, que deve apresentar valores superior a 40 (Reis *et al.*, 2013). O aumento na CP e
713 biomassa total também ocorreu em *B. brizantha* cv. BRS Piatã (Orrico Júnior *et al.*, 2013), cv.
714 Marandu (Pontes *et al.*, 2016) e em *Panicum maximum* (Paciullo *et al.*, 2016) adubadas com
715 nitrogênio.

716 O fósforo (P) é um dos nutrientes mais limitantes para a formação e manutenção da
717 produtividade de pastagens (Dias-Filho, 2011; Mota *et al.*, 2017). Mesmo quando os
718 fertilizantes fosfatados são adicionados aos solos, a oferta de P inorgânico pode sofrer
719 limitação, pois parte desse fósforo pode ser prontamente ligado às partículas do solo (Dias-
720 Filho, 2011; Duchene *et al.*, 2017). No presente estudo, plantas inoculadas apresentaram maior
721 teor de P; o que sugere que as rizobactérias aumentaram a disponibilidade de P no solo. É
722 possível supor, que a inoculação, além de contribuir para o maior desempenho da pastagem,
723 também contribuiria para a melhoria da nutrição animal, diminuindo a necessidade da
724 suplementação de P na dieta animal, o qual é comum em pastagens tropicais (Dias-Filho, 2011).
725 A inoculação com micorriza também promoveu o crescimento em *B. decumbens*, mas não foi
726 capaz de elevar a concentração de P dessa forrageira (Mota *et al.*, 2017).

727 A qualidade da forragem também é determinada pela quantidade de fibras, como a fibra
728 detergente neutra (NDF) e a fibra detergente ácida (ADF), que são carboidratos estruturais que
729 formam a parede celular da planta e incluem celulose, hemicelulose, lignina e pectina. (Ball *et*
730 *al.*, 2015). A qualidade da forragem também está relacionada com a anatomia foliar, pois quanto
731 maior a espessura da epiderme, principalmente a abaxial, e a área de esclerênquima, mais
732 resistente a forrageira será à fermentação dos microrganismos do rumem, diminuindo a
733 digestibilidade (Paciullo *et al.*, 2016; Tsuzukibashi *et al.*, 2016), o que foram menores nas
734 plantas com rizobactérias do que as com adubo. Como forragens com teores de fibras (NDF e
735 ADF) superiores a 60% tem baixa digestibilidade, resultando em menor consumo e desempenho
736 animal (Ball *et al.*, 2015), é possível sugerir que as plantas inoculadas teriam maior potencial
737 de consumo durante o pastejo.

738 No presente estudo, as rizobactérias atuaram como biofertilizantes em *B. brizantha*, por
739 estimular a nutrição, modificar a fisiologia, contribuindo para acúmulo de metabólitos,
740 resultando em maior altura e massa de forragem, o que também seria benéfico em campo,
741 porque aumentaria a massa por bocado e a ingestão diária total de forragem pelos animais (Reis
742 *et al.*, 2013). O uso de rizobactérias aumentou a concentração de N e biomassa em *Brachiaria*
743 spp. (Hungria *et al.*, 2016) e promoveu incremento de nutrientes, produção de biomassa e
744 crescimento em *Triticum aestivum* L. (Rana *et al.*, 2012); *Zea mays* (Dhawi *et al.*, 2015; Armada
745 *et al.*, 2015; Calvo *et al.*, 2017); *Brassica oleracea* (Kumari *et al.*, 2017).

746 Nossos resultados demonstraram que a co-inoculação de *P. fluorescens* and *B.*
747 *pyrrocinia* foi mais eficiente do que a adubação química em *B. brizantha*. A inoculação
748 modificou a anatomia foliar, do colmo e da raiz; incrementou o teor de clorofila, taxas
749 fotossintéticas, eficiência do uso da água, carboidratos totais, amido e proteína; também
750 aumentou os teores de N, P, Na, Mg e Fe nas folhas e raízes, promovendo uma maior qualidade

751 e produção de biomassa em *B. brizantha* (Figura 5). Por tanto, essas rizobactérias poderiam ser
 752 usadas como biofertilizantes em pastos formados por *B. brizantha*, contribuindo para a obtenção
 753 de pastagens de maior produtividade e valor nutritivo.

754 AGRADecIMENTOS

755 Agradecemos a FAPESPA (Fundação de Amparo à Pesquisa do Estado do Pará) pelo
 756 suporte financeiro; Ao Francisco Janyelo Palacios Martinez, do Grupo Gasparim, por fornecer
 757 a semente de *Brachiaria brizantha* cv. BRS Piatã; ao Dr. Walter Velasco e ao Ms. Lenison
 758 Ferreira pelo suporte técnico; e ao Dr. Bruno da Embrapa Agrossilvipastoril pelo suporte nas
 759 análises nutritivas.

760 REFERÊNCIAS

- 761 ARMADA E., AZCON R., LOPEZ-CASTILLO O.M., CALVO-POLANCO M. AND RUIZ-
 762 LOZANO J.M. (2015) Autochthonous arbuscular mycorrhizal fungi and *Bacillus*
 763 *thuringiensis* from a degraded Mediterranean area can be used to improve physiological
 764 traits and performance of a plant of agronomic interest under drought conditions. *Plant*
 765 *Physiology and Biochemistry*, **90**, 64-74.
- 766 BALL D.M., HOVELAND C.S. and G.D. LACEFIELD (2015). Southern Forages. 5th ed.
 767 International Plant Nutrition Institute, Peachtree Corners, GA.
- 768 CALVO P., NELSON L. and KLOEPPER J.W. (2014) Agricultural uses of plant biostimulants.
 769 *Plant Soil*, **383**, 3–41.
- 770 CALVO P., WATTS D.B., KLOEPPER J.W. and TORBERT H.A. (2017) Effect of microbial-
 771 based inoculants on nutrient concentrations and early root morphology of corn (*Zea*
 772 *mays*) *Journal of Plant Nutrition and Soil Science* **180**, 56–70.
- 773 CAPPELLARI L.R., SANTORO M.V., REINOSO H., TRAVAGLIA C., GIORDANO W. and
 774 BANCHIO E. (2015) Anatomical, Morphological, and Phytochemical Effects of
 775 Inoculation with Plant Growth- Promoting Rhizobacteria on Peppermint (*Mentha*
 776 *piperita*). *Journal Chemical Ecology* DOI 10.1007/s10886-015-0549-y
- 777 DIAS-FILHO M.B. (2011) Degradação de pastagens: processos, causas e estratégias de
 778 recuperação. 4ed., Belém. PA.

- 779 DHAWI F., DATTA R. and RAMAKRISHNA W. (2015) Mycorrhiza and PGPB modulate
780 maize biomass, nutrient uptake and metabolic pathways in maize grown in mining-
781 impacted soil. *Plant Physiology and Biochemistry*, **97**, 390-399.
- 782 DUBOIS, M., GILLES, K.A., HAMILTON, J.K., REBERS, P.A. and SMITH, F. (1956)
783 Colorimetric method for determination of sugars and related substances. *Analytical*
784 *Chemistry*. V.28, n.3, p.350-356.
- 785 DUCHENE O., VIAN J.F. and CELETTE F. (2017) Intercropping with legume for
786 agroecological cropping systems: Complementarity and facilitation processes and the
787 importance of soil microorganisms. A review. *Agriculture, Ecosystems and*
788 *Environment*, **240**, 148-161.
- 789 EL-AFRY M.M., EL-NADY M.F., ABDELMONTELEB E.B. and METWALY M.M.S.
790 (2012) Anatomical studies on drought-stressed wheat plants (*Triticum aestivum* L.)
791 treated with some bacterial strains. *Acta Biologica Szegediensis*, **56** (2), 165-174.
- 792 HUNGRIA M., NOGUEIRA M.A. and ARAUJO R.S. (2016) Inoculation of *Brachiaria* spp.
793 with the plant growth-promoting bacterium *Azospirillum brasilense*: An environment-
794 friendly component in the reclamation of degraded pastures in the tropics. *Agriculture,*
795 *Ecosystems & Environment*, **221**, 125-131.
- 796 KADO C. I. and HESKETT M. G. (1970) Selective media for isolation of *Agrobacterium*,
797 *Corynebacterium*, *Erwinia*, *Pseudomonas*, and *Xanthomonas* *Phytopathology*, **60**, 969-
798 976.
- 799 KUMARI S., RAJWADE V., RAMTEKE P., LAWRENCE R. and MASIH H. (2017) Isolation
800 and characterization of potassium and phosphorus solubilising bacteria and fungus
801 (KSB, PSB, KSF, PSF) and its effect on Cauliflower *Int.J.Curr.Microbiol.App.Sci*, **6**(4),
802 987-1006.
- 803 LARRABURU E.E. and LLORENTE B.E. (2015) Anatomical changes induced by
804 *Azospirillum brasilense* in in vitro rooting of pink lapacho. *Plant Cell Tiss Organ Cult*
805 DOI 10.1007/s11240-015-0759-6
- 806 LICHTENTHALER, H.K. (1987). Chlorophyll and carotenoids: pigments of photosynthetic
807 biomembranes. *Methods Enzymology* **148**, 350-382. doi: 10.1016/0076-6879(87)48036-
808 1
- 809 MATHIVANAN S., CHIDAMBARAM A.L.A., ROBERT G.A. and KALAIKANDHAN R.
810 (2017) Impact of PGPR inoculation on photosynthetic pigment and protein contents in
811 *Arachis hypogaea* L. *Journal of Scientific Agriculture*, **1**, 29-36.
- 812 MOTTA P.E.F., SIQUEIRA J.O., RIBEIRO B.T., SILVA S.H.G., POGGERE G.C. and CURI
813 N. (2017) *Urochloa decumbens* growth and P uptake as affected by long-term phosphate
814 fertilization, mycorrhizal inoculation and historical land use in contrasting Oxisols of
815 the Brazilian Cerrado. *Ciência e Agrotecnologia*, **41**(2), 209-219.

- 816 O'BRIEN T.P., FEDER N. and MCCULLY M.E. (1964) Polychromatic staining of plant cell
817 walls by toluidine blue. *Protoplasma*, **59**, 368-373.
- 818 ORRICO JUNIOR M.A.P., ORRICO A.C.A., CENTURION S.R., SUNADA N.S., and LUCAS
819 JUNIOR J. Nutritive value of Piatã grass fertilized with different doses of biofertilizers.
820 *Revista Agrarian*, **6**(21), 312-319.
- 821 PACIULLO, D. S. C., GOMIDE, C. A. M., CASTRO, C. R. T., MAURÍCIO, R. M.,
822 FERNANDES, P. B. AND MORENZ, M. J. F. (2016) Morphogenesis, biomass and
823 nutritive value of *Panicum maximum* under different shade levels and fertilizer nitrogen
824 rates. *Grass and Forage Science*. Version of Record online: 17 OCT 2016, DOI:
825 10.1111/gfs.12264
- 826 PAGANO M.C., CORREA E.J.A., DUARTE N.F., YELIKBAYEV B., O'DONOVAN A. and
827 GUPTA V.K. (2017) Advances in Eco-Efficient Agriculture: The Plant-Soil
828 Mycobiome. *Agriculture*, **7**, 14; doi:10.3390/agriculture7020014
- 829 PII Y., MIMMO T., TOMASI N., TERZANO R., CESCO S., CRECCHIO C. (2015) Microbial
830 interactions in the rhizosphere: beneficial influences of plant growth-promoting
831 rhizobacteria on nutrient acquisition process. A review. *Biol Fertil Soils*. **51**, 403. DOI
832 10.1007/s00374-015-0996-1
- 833 PONTES L.S., BALDISSERA T.C., GIOSTRI A.F., STAFIN G., SANTOS B.R.C. and
834 Carvalho P. C. F. (2016) Effects of nitrogen fertilization and cutting intensity on the
835 agronomic performance of warm-season grasses. *Grass and Forage Science* doi:
836 10.1111/gfs.12267
- 837 PORRA R.J., THOMPSON W.A. and KRIEDEMANN P.E. (1989) Determination of accurate
838 extinction coefficients and simultaneous equations for assaying chlorophylls a and b
839 extracted with four different solvents: Verification of the concentration of chlorophyll
840 standards by atomic absorption spectroscopy. *Biochem Biophys Acta* **975**, 384- 394.
- 841 RANA A., SAHARAN B., NAIN L., PRASANNA R. and SHIVAY Y.S. (2012). Enhancing
842 micronutrient uptake and yield of wheat through bacterial PGPR consortia. *Soil Science*
843 *and Plant Nutrition*, **58**, 573-582.
- 844 REIS A.R., BERNARDES T.F and SIQUEIRA G.R. (2013). Forragicultura: ciência, tecnologia
845 e gestão de recursos forrageiros. 714p.
- 846 RICHARDSON A.E., BAREA J.M., MCNEILL A.M. and C. PRIGENT-COMBARET (2009)
847 Acquisition of phosphorus and nitrogen in the rhizosphere and plant growth promotion
848 by microorganisms. *Plant Soil*, **321**, 305–339. DOI 10.1007/s11104-009-9895-2
- 849 SCHNEIDER, C.A., RASBAND, W.S. and ELICEIRI, K.W. (2012) NIH Image to ImageJ: 25
850 years of image analysis. *Nature Methods* **9**, 671–675. doi:10.1038/nmeth.2089
- 851 STEFAN M., MUNTEANU N., STOLERU V., MIHASAN M. and HRITCU L. (2013) Seed
852 inoculation with plant growth promoting rhizobacteria enhances photosynthesis and
853 yield of runner bean (*Phaseolus coccineus* L.). *Scientia Horticulturae*, **151**, 22–29.

- 854 TAJIMA R. and KATO Y. (2011) Comparison of threshold algorithms for automatic image
855 processing of rice roots using freeware ImageJ. *Field Crops Research* **121**, 460–463.
856 doi:10.1016/j.fcr.2011.01.015
- 857 TIMMUSK S, BEHERS L., MUTHONI J., MURAYA A. and ARONSSON A.C. (2017)
858 Perspectives and challenges of microbial application for crop improvement. *Frontiers*
859 *in Plant Science*, **8**, article 49.
- 860 TSUZUKIBASHI D., COSTA J.P.R., MORO F.V., RUGGIERI A.C. and MALHEIROS E.B.
861 (2016) Quantitative anatomy, in vitro digestibility and chemical composition of
862 *Brachiaria brizantha* cultivars. *Revista de Ciências Agrárias*, **39**(1), 46-53.
- 863 VACHERON J., DESBROSSES G., BOUFFAUD M.L., TOURAINE B., MOËNNE-
864 LOCCOZ Y., MULLER D., LEGENDRE L., WISNIEWSKI-DYÉ F. and C.
865 PRIGENT-COMBARET (2013) Plant growth-promoting rhizobacteria and root system
866 functioning. *Front Plant Science*, **4**, 356. doi: 10.3389/fpls.2013.00356
- 867 VAN SOEST P.J., ROBERTSON J.B. and LEWIS B.A. (1991). Methods for dietary fiber,
868 neutral detergent fiber, and nonstarch polysaccharides in relation to animal production.
869 *J. Dairy Sci.* **74**, 3583-3597.
- 870 WARWATE S.I., KANDOLIYA U.K., BHADJA N.V. and GOLAKIYA B.A. (2017) The
871 effect of plant growth promoting rhizobacteria (pgpr) on biochemical parameters of
872 coriander (*Coriandrum sativum* L.) seedling. *Int.J.Curr.Microbiol.App.Sci.*, **6**(3), 1935-
873 1944.
- 874
- 875
- 876
- 877
- 878
- 879
- 880
- 881
- 882
- 883
- 884

885 Tabelas

886 Tabela 1. Parâmetros anatômicos (μm or μm^2) de *Brachiaria brizantha* não inoculada sem
 887 fertilizante (C-), com fertilizante (C+) e coinoculada com *Pseudomonas fluorescens* (BRM-
 888 32111) e *Burkholderia pyrrocinia* (BRM-32113) (Mix).

Parâmetros		C -	C +	Mix
Folha	Epiderme adaxial	11,54 b	11,86 b	20,24 a
	Epiderme abaxial	16,60 c	25,49 a	23,33 b
	Células buliformes (área)	12828,99 b	12703,61 c	16594,07 a
	Xilema (área)	8100,35 c	8869,52 b	9116,19 a
	Floema (área)	3147,13 a	2160,41 c	2830,56 b
	Esclerênquima (área)	1171,01 b	1472,41 a	800,97 c
	Feixes vasculares (área)	3058,87 b	2686,55 c	3200,26 a
	Bainha dos feixes (área)	18862,34 c	21484,33 b	24316,82 a
	Mesófilo clorofiliano (área)	193524,01 b	188700,81 c	211559,31 a
Colmo	Epiderme	13,47 c	18,46 a	14,07 b
	Feixe vascular (área)	1989,95 c	10089,77 b	20364,01 a
	Número de feixes	33,57 c	42,48 b	69,12 a
Raiz	Epiderme	14508,01 a	13037,02 b	12734,13 c
	Exoderme	18873,81 b	14179,94 c	23063,50 a
	Córtex	209883,81 a	145267,23 c	153069,71 b
	Endoderme	12954,78 c	14006,01 b	14693,67 a
	Periciclo	25278,37 b	23272,28 c	37372,27 a
	Cilindro vascular	338050,4 a	190090,8 b	337155,31 a
	Números de metaxilema	16,52 a	8,24 c	15,30 b
	Números de protoxilema	31,51 b	23,95 c	36,11 a
	Diâmetro do xilema	30041,11 c	38490,77 b	45667,51 a

889 *Significativo ($P < 0.05$, Duncan Test)

890

891

892

893

894

895

896 **Tabela 2** Concentração de macro e micronutrients (g/kg), fibra em detergente neutron (FDN -
 897 g kg⁻¹ of DM), fibra em detergente ácido (FDA - g kg⁻¹ of DM) e proteína bruta (PB - g kg⁻¹ of
 898 DM) de *Brachiaria brizantha* não inoculada sem fertilizante (C-), com fertilizante (C+) e
 899 coinoculada com *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-
 900 32113) (Mix).

Parameters	Folha			Raiz		
	C-	C+	Mix	C-	C+	Mix
N	23.19 c	27.08 b	30.08 a	4.14 c	5.96 b	7.49 a
P	1.07 c	1.17 b	1.22 a	0.07 b	0.08 b	0.17 a
K	24.39 c	23.66 b	22.71 c	8.12 c	9.83 b	14.74 a
Na	0.6 c	0.77 b	0.81 a	1.16 c	1.52 b	1.96 a
Ca	5.94 c	6.88 a	6.79 b	1.12 b	1.53 a	1.09 b
Mg	4.56 c	5.76 b	5.86 a	0.66 c	0.75 b	1.08 a
Fe	191.02 c	212.50 b	542.69 a	2176.9 c	2350.5 b	3143.6 a
Zn	97.12 c	149.13 a	123.37 b	80.86 a	78.98 b	67.03 c
Cu	13.29 c	14.14 b	14.61 a	3.91 c	5.62 a	4.96 b
Mn	143.02 c	169.24 a	145.06 b	53.26 c	61.05 b	74.57 a
FDN	260	251	260			
FDA	130 b	150 a	132 b			
PB	157.6 c	183.3 b	215.8 a			

901 *Significantivo (P < 0.05, Duncan Test).
 902
 903
 904
 905
 906
 907
 908
 909
 910

911 Figuras




Figura 1 Secção transversal de folha (a-c), colmo (d-f) e raiz (g-i) de *Brachiaria brizantha* não inoculada sem fertilizante (a, d, g), com fertilizante (b, e, h) e coinoculada com *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113). 21 dias após a inoculação (35 dias após a emergência das plantas). Célula buliforme (Bc), bainha do feixe (Bs), feixe vascular (Vb), xilema (X), mesófilo (Mes), exoderme (Ex), periciclo (pe), cilindro vascular (vc) e protoxilema (px).




Figura 2 Clorofila (a), índice Spad (b), fotossíntese líquida (A) (c) e eficiência do uso da água (WUE) de *Brachiaria brizantha* não inoculada sem fertilizante (C-), com fertilizante (C+) e coinoculada com *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113) (Mix). Médias seguidas da mesma letra na coluna não diferem ($P < 0,05$, teste de Duncan).




Figura 3 Amido (a) e carboidratos totais (b) de *Brachiaria brizantha* não inoculada sem fertilizante (C-), com fertilizante (C+) e coinoculada com *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113) (Mix). Médias seguidas da mesma letra em cada coluna não diferem ($P < 0,05$, teste de Duncan).





Figura 4 Altura (H) (a), área radicular (b) e produção de biomassa (c) de *Brachiaria brizantha* não inoculada sem fertilizante (C-), com fertilizante (C+) e coinoculada com *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113) (Mix). Médias seguidas da mesma letra na coluna não diferem ($P < 0,05$, teste de Duncan).



948

949 **Figura 5** *Brachiaria brizantha* coinoculada com rizobactérias promotoras de crescimento em
 950 plantas (PGPR), 21 dias após a inoculação (35 dias após a emergência das plantas).
 951 *Pseudomonas fluorescens* (BRM-32111) e *Burkholderia pyrrocinia* (BRM-32113)
 952 modificaram a anatomia (folha, colmo e raiz), incrementaram a altura (H), taxa fotossintética
 953 (*A*), eficiência do uso da água (WUE), concentração de clorofila, N, P, Na, Mg, Fe, amido,
 954 carboidrato, proteína, área radicular, aumentando a qualidade de forragem e produção de
 955 biomassa em *B. brizantha*.

956

957

958

959

960

961

962 **Light and plant growth-promoting rhizobacteria (PGPR) effects on *Brachiaria brizantha***
 963 **growth and phenotypic plasticity to shade³**

964 Monyck Jeane dos Santos Lopes, Moacyr Bernardino Dias Filho, Thomaz Henrique dos Reis
 965 Castro, Gisele Barata da Silva

966 **Abstract**

967 This is the first report on the effect of light intensity and plant growth-promoting
 968 rhizobacteria (PGPR) on the growth of a tropical forage grass, being a relevant study to improve
 969 pasture management in conventional farming and integrated crop-livestock-forestry systems.
 970 In this study, our aim was to evaluate the effects of light intensity and *Burkholderia pyrrocinia*
 971 and *Pseudomonas fluorescens* inoculation on *Brachiaria brizantha* cv. BRS Piatã growth,
 972 phenotypic plasticity and shade tolerance. The experiment was conducted in a semi-controlled
 973 environment. Seedlings of *B. brizantha* were allocated to full sun and shade. *P. fluorescens* and
 974 *B. pyrrocinia* were inoculated individually or co-inoculated by soil drench, 14 days after
 975 seedling emergence. We evaluated morphogenesis, structural and growth parameters.
 976 Irrespective of the light regime, co-inoculated plants had greater leaf area and SPAD index
 977 (chlorophyll content). Increase in total biomass production in co-inoculated plants was over
 978 100% and 300%, under full sun and shade, respectively. Co-inoculated *P. fluorescens* and *B.*
 979 *pyrrocinia* increased shade tolerance in *B. brizantha*, improving plant performance. Co-
 980 inoculation promoted growth in *B. brizantha* under both sun and shade, indicating its potential
 981 as a bio-fertilizer in conventional and integrated systems, especially in silvipastoral systems,
 982 where light availability to pasture growth may be limited.

983 **Keywords:** shade, forage grass, *Pseudomonas fluorescens*, *Burkholderia pyrrocinia*

³ Este capítulo segue as normas de formatação da **Grass and Forage Science**.

984 INTRODUCTION

985 Integrated crop-livestock-forestry systems are efficient land-management alternatives
 986 for restoring degraded pastures, increasing food security and promoting carbon sequestration
 987 (Dias-Filho, 2011; Moraes *et al.*, 2014; Paciullo *et al.*, 2016; Santos *et al.*, 2016). However, as
 988 the forage grass species planted in integrated systems may be subjected to light restriction,
 989 imposed by agricultural crops or tree species, it is necessary to know their agronomic
 990 performance under shade, to define their potential of use in these systems (Dias-Filho, 2000;
 991 Paciullo *et al.*, 2011; Gomez *et al.*, 2012; Pimentel *et al.*, 2016).

992 Phenotypic plasticity relates to the species adaptability to changes in the environment,
 993 such as variations in solar radiation, temperature, soil water and nutrient availability (Gomez *et*
 994 *al.*, 2012; Valladares *et al.*, 2016). Shade tolerant plants can adjust their morphophysiology,
 995 such as biomass allocation pattern, chlorophyll content and leaf area and thickness to maximize
 996 light capture (Paciullo *et al.*, 2016; Valladares *et al.*, 2016)

997 In conventional and integrated systems, forage grass production is usually limited by
 998 low soil fertility (Paciullo *et al.*, 2011; Paciullo *et al.*, 2016; Pimentel *et al.*, 2016). The use of
 999 plant-growth-promoting rhizobacteria (PGPR) could be an alternative for reducing the use of
 1000 chemical fertilizers, with clear environmental and economic benefits (Nadeem *et al.*, 2014;
 1001 Paredes and Lebeis, 2016). The beneficial effect of PGPR on forage grasses has been reported
 1002 for *Azospirillum brasilense* on *B. briazantha* (Hungria *et al.* 2016), and *Azotobacter*,
 1003 *Azospirillum* and *Herbaspirillum* on *Axonopus affinis*, *Paspalum notatum*, *Andropogon*
 1004 *lateralis* and *Aristida laevis* (Marques *et al.*, 2017).

1005 Beneficial microorganisms are known to improve nutrient uptake, phosphorus
 1006 solubilization, phytohormone production and disease resistance by elicited induced systemic
 1007 resistance or systemic acquired resistance. Also, beneficial microorganisms modify the

phenotypic plasticity of plants, by mitigating the negative impact of abiotic stresses (Goh *et al.*, 2013; Vacheron *et al.*, 2013; Paredes and Lebeis, 2016; Vimal *et al.*, 2017), including light limitation (Konvalinková and Jansa, 2016).

Previous studies with *Pseudomonas* sp. and *Burkholderia* sp. attested their growth promote potential in rice (Rêgo *et al.*, 2014; Nascente *et al.*, 2016) and *Brachiaria brizantha* (exploratory research conducted by the first author at the Federal Rural University of Amazonia). These rhizobacteria are known to increase the auxin synthesis, nutrient uptake (e.g., nitrogen, phosphorus, and iron), chlorophyll content, photosynthetic rate, and biomass production (Ahemad and Kibret, 2014; Nascente *et al.*, 2016).

Although light is known to interfere in mycorrhiza-plant mutualism (Saner *et al.*, 2011; Aguilar-Chama and Guevara, 2016; Konvalinková and Jansa, 2016), we could not find any published study on the effect of light intensity and use of rhizobacteria on the growth and phenotypic plasticity of a tropical forage grass species.

In Brazil, *Brachiaria* is the most important grass genus for pasture formation (Reis *et al.*, 2013; Santana *et al.*, 2016; Santos *et al.*, 2016), which stands out *B. brizantha* (Hochst ex. A. Rich.) cv. BRS Piatã (Reis *et al.*, 2013). We hypothesized that *Burkholderia pyrrocinia* and *Pseudomonas fluorescens* inoculation can increase *B. brizantha* growth under limited light conditions, by increasing the shade tolerance of this grass cultivar. In this study, our aim was to evaluate the effects of light intensity and *B. pyrrocinia* and *P. fluorescens* inoculation on the growth, phenotypic plasticity and shade tolerance of *B. brizantha* cv. BRS Piatã.

MATERIALS AND METHODS

Study site, plant material and inoculant

1030 The experiment was conducted in a semi-controlled environment in the nursery seedling
 1031 production unit of the Federal Rural University of Amazonia (UFRA) (01°27'25 "S, 48°26'36"
 1032 W) in Belém, Pará, Brazil. The regional climate here according to Koppen classification is Af
 1033 (equatorial). During the experimental period the mean air temperature and relative humidity
 1034 were $32 \pm 2,8^{\circ}\text{C}$ and $73 \pm 3\%$ (mean $\pm$ s.d.), respectively.

1035 Seeds of *B. brizantha* cv. BRS Piatã were sown in polyethylene pots (15 x 25 x 0.05
 1036 cm) filled with soil (Ferralsol - top soil from a second growth forest - pH, 4,2; organic
 1037 matter, $18,80 \text{ g dm}^{-3}$; P, 2 mg dm^{-3} ; K, 4 mg dm^{-3} ; Ca, $0,2 \text{ mmolc dm}^{-3}$; Ca + Mg, $0,3 \text{ mmolc}$
 1038 dm^{-3} ; Al, $1,4 \text{ mmolc dm}^{-3}$) and kept under greenhouse conditions. Plants were grown under full-
 1039 sun and artificial shade (0% and 47% of shade, or $2100 \mu\text{molm}^{-2} \text{ s}^{-1}$ and $1113 \mu\text{molm}^{-2} \text{ s}^{-1}$, at
 1040 canopy height, measured at 11 am, local time, on a cloudless day) obtained with black
 1041 polyethylene screens (solar radiation transmissivity of ca. 50%) placed 160 cm above the
 1042 canopy height.

1043 We used *P. fluorescens* (BRM-32111) and *B. pyrrocinia* (BRM-32113), supplied by the
 1044 Plant Protection Laboratory of the Federal Rural University of Amazonia *in vitro* collection,
 1045 and originally selected from the rhizosphere of rice plants. The rhizobacteria were cultured in
 1046 Petri dishes with a solid culture medium 523 (agar, casein hydrolysate, magnesium sulfate
 1047 anhydrous, potassium phosphate monobasic, sucrose and yeast extract) (Kado and Heskett,
 1048 1970) and incubated for 48h at 28°C . The bacterial suspension was prepared in sterile water and
 1049 adjusted to 550 nm (10^8 CFU). Soil-drenched inoculation was carried out 14 days after seedling
 1050 emergence, each pot received a 5mL bacterial suspension. The experiment was repeated three
 1051 times with similar results.

1052 Plant growth parameters

1053 Morphogenetic and structural parameters calculated according to Gomide and Gomide
 1054 (2000), were: leaf appearance rate (ratio between the difference in the number of initial and
 1055 final leaves and the number of days of the evaluation interval days, LApR), leaf elongation rate
 1056 (ratio between the difference of the initial and final lengths of the expanded laminae and the
 1057 number of days of the evaluation interval, LER), culm elongation rate (ratio between the
 1058 difference of the initial and final lengths of culm and the number of days of the evaluation
 1059 interval, CER), number of leaves per plant (NL) and number of tillers per plant (NT). We also
 1060 determined the plant height (H), culm length (CL) and we measured the chlorophyll content
 1061 (SPAD index - soil plant analysis development) in the youngest fully expanded leaf blade. Each
 1062 SPAD index value was the mean of five readings per leaf. We used a portable chlorophyll meter
 1063 (SPAD-502. Konica Minolta Sensing, INC. Japan).

1064 At 35 days after seedling emergence, five plants per treatment were harvested and
 1065 separated into shoot (leaf blades and culms) and roots. Plant material was oven dried (60° C)
 1066 until constant mass. Total dry mass (TDM) was calculated by adding shoot dry mass (SDM)
 1067 and root dry mass (RDM). The biomass allocation pattern was estimated as the leaf, culm and
 1068 root mass ratios (respectively, the ratio between total leaf, culm, and root dry mass per plant
 1069 and total dry mass per plant).

1070 Specific leaf area (SLA, the ratio of leaf area to leaf dry mass) was determine over leaf
 1071 disks of either 0.42 cm² or 2.28 cm², dried at 60°C until constant mass. Total leaf area per plant
 1072 was estimated from SLA and leaf mass results. We also calculated root dry mass/shoot dry mass
 1073 ratio (RDM/SDM) and leaf area ratio (ratio of leaf area per total dry mass per plant, LAR).
 1074 Relative growth rate (change in total mass per total dry mass of plant per day, RGR) was
 1075 calculated for harvests at 14 and 35 days after seedling emergence. The allometric coefficient

1076 K was calculated as the ratio of the growth rate of root and shoot. All plant growth parameters
 1077 were calculated according to Hunt (1990) and Barbero *et al.* (2013).

1078 Statistical analyses

1079 The experimental design was completely randomized in a 4×2 factorial arrangements
 1080 (non-inoculated *B. brizantha*, inoculated with BRM-32111, inoculated with BRM-32113 and
 1081 co-inoculated (BRM-32111 + BRM-32113) x full-sun and shade), with five replicates. Data were
 1082 subjected to analysis of variance and, when appropriate, the means were compared by Duncan
 1083 test at 5%. The assumption of homogeneity of variances and normality were tested for each
 1084 ANOVA and, when necessary, data were log, or square root transformed. Transformed values
 1085 were back transformed for presentation. Statistical analyses were performed by the statistical
 1086 package STATISTICA (StatSoft, Inc., Tulsa, USA).

1087 RESULTS

1088 When the bacteria were inoculated individually, plants under full-sun showed the
 1089 highest growth with *P. fluorescens* (BRM-32111), and with *B. pyrrocinia* (BRM-32113), when
 1090 grown under shade (Figure 1). The maximum increase in height ($F_{3,32} = 38.05$; $P < 0.001$)
 1091 resulted from the activity of the BRM-32113, under full-sun (18%), and of BRM-32111, under
 1092 shade (13%) (Figure 2). Leaf appearance ($F_{3,32} = 11.64$; $P < 0.001$) and elongation ($F_{3,32} =$
 1093 57.14 ; $P < 0.001$), culm elongation ($F_{3,32} = 24.97$; $P < 0.001$), number of leaves ($F_{3,32} = 43.02$;
 1094 $P < 0.001$) and leaf length ($F_{3,32} = 5.98$; $P < 0.05$), were higher for BRM-32113, in full-sun
 1095 plants, and, in most cases, for BRM-32111, in shaded plants (Table 1).

1096 Higher values of specific leaf area (SLA) ($F_{3,32} = 245.64$; $P < 0.001$), and leaf area ratio
 1097 (LAR) ($F_{3,32} = 347.52$; $P < 0.001$), were obtained in non-inoculated plants, in the shade. Under
 1098 full sun, the highest SLA and LAR values were found in plants inoculated with BRM-32113
 1099 (Figure 3). Leaf area increased by 122% in full sun plants, inoculated with BRM-32113 and by

1100 20% in shade plants, inoculated with BRM-32111 (Figure 4). When inoculated individually,
 1101 the SPAD index increased ($F_{3,32} = 6.18$; $P < 0.05$) by 4%, in full sun plants and by 7%, in shade
 1102 plants (Figure 4).

1103 When the rhizobacteria were inoculated individually, biomass production ($F_{3,32} =$
 1104 616.61 ; $P < 0.001$) and relative growth rate ($F_{3,32} = 119.97$; $P < 0.001$) were increased in full-
 1105 sun plants by 30% (BRM-32113) and, under shade, by more than 100% (BRM-32111) (Table
 1106 2; Figure 5). Under shade, inoculation favored an increased biomass allocation to roots (Figure
 1107 6). Under full sun, the root to shoot biomass ratio ($F_{3,32} = 42.71$; $P < 0.05$) and allometric index
 1108 (K) ($F_{3,32} = 57.16$; $P < 0.0001$) were higher in non-inoculated plants and in plants inoculated
 1109 with BRM-32111. The reason for that was a preferential biomass allocation to the roots, while
 1110 in plants inoculated with BRM-32111, biomass was allocated mainly to leaves (Table 2; Figure
 1111 6).

1112 Co-inoculation promoted growth of *B. brizantha* under both full-sun and in the shade
 1113 (Figure 1). Plant height under co-inoculation increased by 28% and 65%, under full-sun and
 1114 shade, respectively, relative to non-inoculated plants (Figure 2). Higher leaf appearance and
 1115 elongation rates, culm elongation, number of leaves and leaf length occurred in co-inoculated
 1116 plants, regardless of light regime (Table 1).

1117 Tillering was observed only in plants grown under full sun, being 50% higher in co-
 1118 inoculated plants (Table 1). The increase in the culm length and in the leaf and culm elongation
 1119 rates, in response to shade, was higher in co-inoculated plants (Table 1). The co-inoculated
 1120 plants had a 130% larger leaf area, in full sun plants and a 200% larger leaf area in shade plants
 1121 (Figure 4). The SPAD index was higher in co-inoculated plants, increasing by 13% in full sun
 1122 plants and by 17% in shade plants.

1123 Co-inoculation increased biomass production, irrespective of the light intensity (Fig. 5).
 1124 Under full sun co-inoculation increased leaf (139%), culm (86%) and root (88%) biomass,
 1125 resulting in an increase of over 100%, in total biomass relative to non-inoculated plants (Table
 1126 2; Fig. 5). Under shade, co-inoculation increased leaf (300%), culm (500%) and root (670%)
 1127 biomass, resulting in an increase of about 300%, in total biomass, relative to non-inoculated
 1128 plants (Table 2; Fig. 5).

1129 Co-inoculation also increased relative growth rate by 110%, under full sun, and 500%,
 1130 in shaded plants (Figure 5). Shaded plants showed higher values of root to shoot biomass ratio
 1131 and allometric index (K), particularly when co-inoculated (Table 2). Under full sun, co-
 1132 inoculated plants allocated proportional amounts of biomass between leaves and roots (Fig. 6).

1133 DISCUSSION

1134 When inoculation was performed individually, the benefit varied with the type of
 1135 bacteria and light intensity. Growth of plants inoculated only with *B. pyrrocinia* was positively
 1136 related to light intensity. A possible cause for this response was a likely higher demand in
 1137 photoassimilates for the symbiosis of this microorganism (Aguilar-Chama and Guevara, 2016;
 1138 Konvalinková and Jansa, 2016). A contrasting response occurred in plants inoculated only with
 1139 *P. fluorescens*, which fostered higher growth mainly under restricted light intensities.

1140 It could be inferred that the higher radiation intensity might have modified the quantity
 1141 and chemistry composition of root-exudates, affecting rhizodeposition processes and disturbing
 1142 rhizosphere functioning (Haichar *et al.*, 2008; Venturi and Keel, 2016; Vimal *et al.*, 2017). This
 1143 could have interfered in the quorum sensing, inhibiting the mutual interaction under full sun
 1144 (Goh *et al.*, 2013; Venturi and Keel, 2016). This response supports the concept that light can
 1145 interfere with mutualistic interaction, which in turn, varies per the host, and beneficial
 1146 microorganisms (Konvalinková and Jansa, 2016). Under limited light conditions, microbial root

1147 symbionts can create additive costs, resulting in decreased plant fitness, as observed in *Vatica*
 1148 *albiramis* (Saner *et al.*, 2011) and in *Datura stramonium* (Aguilar-Chama and Guevara, 2016),
 1149 where growth promotion is positively related to light intensity, occurring only at higher light.

1150 When co-inoculated, *B. pyrrocinia* and *P. fluorescens* promoted growth of *B. brizantha*,
 1151 irrespective of light intensity. The rhizosphere microbiota is known to extend the adaptive
 1152 capacity of plants to environmental stresses (Venturi and Keel, 2016). The overall beneficial
 1153 effect of the interaction between microorganisms and plant growth, under contrasting light
 1154 environments, has also been reported in studies with mycorrhizal growth promoters (Casierra-
 1155 Posada *et al.*, 2013; Aguilar-Chama and Guevara, 2016; Konvalinková and Jansa, 2016).

1156 In rice, from which the rhizobacteria used in this study were originally isolated, growth
 1157 is higher with *B. pyrrocinia* inoculation than with *P. fluorescens* inoculation (Rêgo *et al.*, 2014;
 1158 Nascente *et al.*, 2016), or with co-inoculation with these rhizobacteria (Rêgo *et al.*, 2014). The
 1159 synergistic effect of *B. pyrrocinia* and *P. fluorescens*, promoting growth in *B. brizantha*, was
 1160 possibly a response of increased nutrient flow (Casierra-Posada *et al.*, 2013, Paredes and
 1161 Lebeis, 2016), phytohormone production, such as auxin or cytokinin, or decreased plant
 1162 ethylene levels, through the action of the enzyme ACC deaminase (Ahemad and Kibret, 2014;
 1163 Vimal *et al.*, 2017).

1164 It can be inferred that co-inoculation increased phenotypic plasticity in various key traits
 1165 as a strategy to mitigate the effect of shade in *B. brizantha*. This was achieved by increasing
 1166 plant height, culm and leaf elongation (i.e., etiolation) (Goh *et al.*, 2013; Martins *et al.*, 2014;
 1167 Valladares *et al.*, 2016), hence increasing total plant biomass. Since tree shading (Santos *et al.*,
 1168 2016) and crop shading (Neves Neto *et al.*, 2015) are known to reduce forage production in *B.*
 1169 *brizantha* cv. BRS Piatã pastures, it could be inferred that co-inoculation would be a suitable
 1170 strategy to counteract limited pasture growth in integrated systems.

Specific leaf area (SLA) and leaf area ratio (LAR) vary with light, temperature, moisture and nutrient availability (Barbero *et al.*, 2013; Daniagry and Dang, 2014). Increased SLA and LAR under low light is a strategy to improve light capture in *B. brizantha* (Dias-Filho, 2000), and this response was observed in non-inoculated plants in our study. Our results also indicate that the ability of *B. brizantha* to modify SLA and LAR under low light can be influenced by plant growth-promoting rhizobacteria. A possible cause for this response is the potential ability of the rhizobacteria to supply the needed soil nutrient resources for shaded *B. brizantha*. This is attained through increased root growth, making it unnecessary for the shaded plants to increase SLA and LAR to those levels observed in uninoculated plants. This indirect effect could make inoculated plants more tolerant to additional stresses such as grazing, increasing pasture performance and productivity (Pimentel *et al.*, 2016).

Considering that the greater the SPAD index, the higher the chlorophyll and nitrogen content of *B. brizantha* leaves (Martucello *et al.*, 2009), presumably co-inoculated rhizobacteria also acted synergistically to increase the nitrogen content of leaves. This is because plant growth-promoting rhizobacteria can also affect plant nutrient-acquisition processes, influencing nutrient availability in the rhizosphere, improving the plant nutritional status, thus increasing pasture growth and productivity (Pii *et al.*, 2015). Higher SPAD values correlating positively with chlorophyll and nitrogen contents, were also found in *B. brizantha* with higher levels of nitrogen fertilization (Lima *et al.*, 2016).

The general increase in root biomass under co-inoculation and the biomass allocation patterns to full-sun and shade plants observed in this study, probably increased the efficiency of use of water and nutrients. The observed responses of co-inoculated plants would be useful to increase the fitness of *B. brizantha* in water- limited environments. In grazed pastures, resources use efficiency ensures the persistence of plants over time and longevity of pastures

(Dias Filho, 2000; Martucello *et al.*, 2009; Pimentel *et al.*, 2016). Also, the increased relative growth rate, tillering, leaf area and biomass production in co-inoculated plants, would be a desirable trait in intensively managed systems, such as rotational grazing systems, reducing grazing intervals.

In conclusion, our results attest the potential of co-inoculated *P. fluorescens* and *B. pyrrocinia* to increase biomass production in *B. brizantha* and the expression of those plant characters that may enhance persistence under reduced light availability. In addition, these findings set up the basis for additional exploratory studies, particularly on the ability of this beneficial interaction to remain fully active under grazing, in integrated crop-livestock-forestry systems.

ACKNOWLEDGEMENTS

We thank FAPESPA (Fundação de Amparo à Pesquisa do Estado do Pará) for financial support. We also thank Francisco Janyelo Palacios Martinez, from Grupo Gasparim, for kindly supplying the *Brachiaria brizantha* cv. BRS Piatã seeds, and Walter Velasco and Lenison Ferreira for technical laboratory support.

No conflict of interest

REFERENCES

- AGUILAR-CHAMA A. and GUEVARA R. (2016) Resource allocation in an annual herb: Effects of light, mycorrhizal fungi, and defoliation. *Acta Oecologica*, **71**, 1-7.
- AHEMAD M. and KIBRET M. (2014) Mechanisms and applications of plant growth promoting rhizobacteria: Current perspective. *Journal of King Saud University – Science*, **26**, 1–20.
- BARBERO L.M., PRADO T.F., BASSO K.C., LIMA L.A., MOTTA K.M., KRÜGER B.C., MARTINS NETO L.R. and SILVA G.A.S. (2013) Análise de crescimento em plantas forrageiras aplicada ao manejo de pastagens. *Veterinária Notícias*, **19**, 71-85.
- CASIERRA-POSADA F., PEÑA-OLMOS J., PEÑALOZA J. and ROVEDA G. (2013) Influencia de la sombra y de las micorrizas sobre el crecimiento de plantas de lulo (*Solanum quitoense* Lam.). *Revista U.D.C.A Actualidad & Divulgación Científica*, **16**, 61–70.

- 1224 DIAS-FILHO M.B. (2000) Growth and biomass allocation of the C4 grasses *Brachiaria*
1225 *brizantha* and *B. humidicola* under shade. *Pesquisa agropecuária brasileira*, **35**, 2335-
1226 2341.
- 1227 DIAS-FILHO M.B. (2011) Degradação de pastagens: processos, causas e estratégias de
1228 recuperação. 4ed., Belém. PA.
- 1229 GOH C.H., VALLEJOS D.F.V., NICOTRA A.B. and MATHESIUS U. (2013) The impact of
1230 beneficial plant-associated microbes on plant phenotypic plasticity. *Journal of*
1231 *Chemical Ecology*, **39**, 826–839.
- 1232 GOMEZ S., GUENNI O. AND BRAVO DE GUENNI L. (2012) Growth, leaf photosynthesis
1233 and canopy light use efficiency under differing irradiance and soil N supplies in the
1234 forage grass *Brachiaria decumbens* Stapf. *Grass and Forage Science*, **68**, 395–407.
- 1235 GOMIDE C.A.M. and GOMIDE J.A. (2000) Morfogênese de cultivares de *Panicum maximum*
1236 Jacq. *Revista Brasileira de Zootecnia*, **29**, 341-348.
- 1237 HAICHAR F.Z., MAROLI C., BERGE O., RANGEL-CASTRO J.I., PROSSER J.I.,
1238 BALESDENT J., HEULIN T. AND ACHOUAK W. (2008) Plant host habitat and root
1239 exudates shape soil bacterial community structure. *The ISME Journal*, **2**, 1221–1230.
- 1240 HUNGRIA M., NOGUEIRA M.A. and ARAUJO R.S. (2016) Inoculation of *Brachiaria* spp.
1241 with the plant growth-promoting bacterium *Azospirillum brasilense*: An environment-
1242 friendly component in the reclamation of degraded pastures in the tropics. *Agriculture,*
1243 *Ecosystems & Environment*, **221**, 125–131.
- 1244 HUNT R. (1990) Basic growth analysis for beginners, London.
- 1245 KADO C. I. and HESKETT M. G. (1970) Selective media for isolation of *Agrobacterium*,
1246 *Corynebacterium*, *Erwinia*, *Pseudomonas*, and *Xanthomonas* *Phytopathology*, **60**, 969–
1247 976.
- 1248 KONVALINKOVÁ T. and JANSÁ J. (2016) Lights off for arbuscular mycorrhiza: on its
1249 symbiotic functioning under light deprivation. *Frontiers in Plant Science*, **7**, 782.
- 1250 LIMA J.E.S., NASCENTE A.S., LEANDRO W.M. and SILVEIRA P.M. (2016) *Urochloa*
1251 *ruziziensis* responses to sources and doses of urea. *Revista brasileira de engenharia*
1252 *agrícola ambiental*, **20**, 401-407.
- 1253 MARQUES A.C.R., OLIVEIRA L.B., NICOLOSO F.T., JACQUES R.J.S., GIACOMINI S.J.
1254 and QUADROS F.L.F. (2017) Biological nitrogen fixation in C4 grasses of different
1255 growth strategies of South America natural grasslands *Applied Soil Ecology*, **113**, 54–
1256 62.
- 1257 MARTINS A.D., SOUSA L.F., NÓBREGA E.B., DONIZETTI J.G.S, SANTOS A.C. and
1258 SOUSA J.T.L. (2014) Relação do nível de sombreamento artificial e da adubação sobre
1259 o desenvolvimento da forrageira *Urochloa brizantha* cv. Marandu. *Revista Brasileira*
1260 *de Saúde e Produção Animal*, **15**, 994-1005.
- 1261 MARTUSCELLO J.A., JANK L., GONTIJO NETO M.M., LAURA V.A. and CUNHA
1262 D.N.F.V. (2009) Produção de gramíneas do gênero *Brachiaria* sob níveis de
1263 sombreamento. *Revista Brasileira de Zootecnia*, **38**, 1183-1190.
- 1264 MORAES A., CARVALHO P.C.F., LUSTOSA S.B.C., LANG C.R. and DEISS L. (2014)
1265 Research on Integrated Crop-Livestock Systems in Brazil. *Revista de Ciência*
1266 *Agronomica*, **45**, 1024-1031.
- 1267 NADEEM S.M., AHMAD M., ZAHIR Z.A., JAVAID A. and ASHRAF M. (2014) The role of
1268 mycorrhizae and plant growth promoting rhizobacteria (PGPR) in improving crop
1269 productivity under stressful environments. *Biotechnology Advances*, **32**, 429–448.
- 1270 NASCENTE A.S., FILIPPI M.C.C., LANNA A.C., SOUZA A.C.A., LOBO V.L.S. and SILVA
1271 G.B. (2016) Biomass, gas exchange, and nutrient contents in upland rice plants affected

- 1272 by application forms of microorganism growth promoters. *Environmental Science and*
 1273 *Pollution Research*
- 1274 NEVES NETO D.N., SANTOS A.C., ALEXANDRINO E. and SANTOS P.M. (2015)
 1275 Características morfológicas e estruturais de *Urochloa* spp. sob manejo convencional e
 1276 consorciado com cereais. *Revista Ciência Agronômica*, **46**, 204-215.
- 1277 PACIULLO D.S.C., FERNANDES P.B., GOMIDE C.A.M., CASTRO C.R.T., SOBRINHO
 1278 F.S. and CARVALHO C.A.B. (2011) The growth dynamics in *Brachiaria* species
 1279 according to nitrogen dose and shade. *Revista Brasileira de Zootecnia*, **40**, 270-276.
- 1280 PACIULLO, D. S. C., GOMIDE, C. A. M., CASTRO, C. R. T., MAURÍCIO, R. M.,
 1281 FERNANDES, P. B. AND MORENZ, M. J. F. (2016) Morphogenesis, biomass and
 1282 nutritive value of *Panicum maximum* under different shade levels and fertilizer nitrogen
 1283 rates. *Grass and Forage Science*. Version of Record online: 17 OCT 2016,
 1284 DOI: 10.1111/gfs.12264
- 1285 PAREDES S.H. and LEBEIS S.L. (2016) Giving back to the community: microbial
 1286 mechanisms of plant–soil interactions. *Functional Ecology*, **30**, 1043–1052.
- 1287 PII Y., MIMMO T., TOMASI N., TERZANO R., CESCO S. and CRECCHIO C. (2015)
 1288 Microbial interactions in the rhizosphere: beneficial influences of plant growth-
 1289 promoting rhizobacteria on nutrient acquisition process. A review. *Biology and Fertility*
 1290 *of Soils*, **5**, 403.
- 1291 PIMENTEL R.M., BAYÃO G.F.V., LELIS D.L., CARDOSO A.J.S., SALDARRIAGA F.V.,
 1292 MELO C.C.V., SOUZA F.B.M., PIMENTEL A.C.S., FONSECA D.M. and SANTOS
 1293 M.E.R. (2016) Ecofisiologia de plantas forrageiras. *PUBVET* **10**, 666-679.
- 1294 REIS A.R., BERNARDES T.F and SIQUEIRA G.R. (2013). Forragicultura: ciência, tecnologia
 1295 e gestão de recursos forrageiros. 714p.
- 1296 RÊGO M.C.F., BORGES F.I., FILIPPI M.C.C., GONÇALVES L.A. and SILVA G.B. (2014)
 1297 Morphoanatomical and biochemical changes in the roots of rice plants induced by plant
 1298 growth-promoting microorganisms. *Journal of Botany*.
- 1299 SANER P., PHILIPSON C., ONG R.C., MAJALAP N., EGLI S. and HECTOR A. (2011)
 1300 Positive effects of ectomycorrhizal colonization on growth of seedlings of a tropical tree
 1301 across a range of forest floor light conditions. *Plant Soil*, **338**, 411–421.
- 1302 SANTANA S.S., BRITO L.F., AZENHA M.V., OLIVEIRA A.A., MALHEIROS E.B.,
 1303 RUGGIERI A.C. and REIS R.A. (2016) Canopy characteristics and tillering dynamics
 1304 of Marandu palisade grass pastures in the rainy–dry transition season *Grass and forage*
 1305 *Science*. Version of Record online: 21 JUN 2016, DOI: 10.1111/gfs.12234.
- 1306 SANTOS D., GUIMARÃES JÚNIOR R., VILELA L., PULROLNIK K., BUFON V.B. and
 1307 FRANÇA A.F.S. (2016) Forage dry mass accumulation and structural characteristics of
 1308 Piatã grass in silvopastoral systems in the Brazilian savannah. *Agriculture, Ecosystems*
 1309 *& Environment*, **233**, 16-24.
- 1310 VACHERON J., DESBROSSES G., BOUFFAUD M.L., TOURAINE B., MOËNNE-
 1311 LOCCOZ Y., MULLER D., LEGENDRE L., WISNIEWSKI-DYÉ F. and PRIGENT-
 1312 COMBARET C. (2013) Plant growth-promoting rhizobacteria and root system
 1313 functioning. *Frontiers in Plant Science*, **4**, 1-19.
- 1314 VALLADARES F., LAANISTO L., NIINEMETS Ü. and ZAVALA M.A. (2016): Shedding
 1315 light on shade: ecological perspectives of understorey plant life. *Plant Ecology &*
 1316 *Diversity* 1-15.
- 1317 VENTURI V. and KEEL C. (2016) Signaling in the Rhizosphere. *Trends in Plant Science* **21**,
 1318 187–198.
- 1319 VIMAL S.R., SINGH J.S., ARORA N.K. and SINGH S. (2017) Soil-Plant-Microbe
 1320 Interactions in Stressed Agriculture Management: A Review. *Pedosphere*, **27**, 177–192.

1321 TABLES

1322 **Table 1** Light intensity and plant growth-promoting rhizobacteria (PGPR) on number of tillers
 1323 (NT), number of leaves per plant (NL), expanded leaf length (EF - cm), culm length (C - cm),
 1324 leaf appearance rate (LApR - L⁻¹.day⁻¹), leaf elongation rate (LER - L⁻¹.day⁻¹) and culm
 1325 elongation rate (CER - L⁻¹.day⁻¹) of *Brachiaria brizantha*

TRATAMENTS		NT	NL	EF	C	LApR	LER	CER
Full sun	Non-inoculated	2 c	11 d	24.83 d	10.80 c	0.37 c	0.88c	0.43 b
	BRM-32111	2 c	13 c	27.88 c	11.75 b	0.45 b	0.84d	0.42 b
	BRM - 32113	2 b	16 b	33.6 b	12.05 a	0.45 b	0.96 b	0.44 b
	MIX	3 a	17 a	36.5 a	12.01 a	0.51 a	1.32 a	0.60 a
Shade	Non-inoculated	0	5 d	21.91 c	10.35 d	0.10 c	0.61c	0.26 d
	BRM - 32111	0	6 b	27.77 b	13.83 b	0.20 b	1.45b	0.63 b
	BRM - 32113	0	5 c	28.2 b	11.66 c	0.18 b	1.41 b	0.56 c
	MIX	0	6 a	34.07 a	15.60 a	0.3 a	2.05 a	0.88 a

1326 Means followed by different letters in each column and within each light intensity are
 1327 significantly different (P < 0.05, Duncan Test). BRM-32111 = *Pseudomonas fluorescens*;
 1328 BRM-32113 = *Burkholderia pyrrocinia*; MIX= BRM-32111 + BRM-32113.

1329

1330 **Table 2** Light intensity and plant growth-promoting rhizobacteria (PGPR) on the leaf (LDM -
 1331 g), culm (CDM - g), root (RDM - g) and root/shoot dry mass production, and allometric
 1332 coefficient K (growth rate of root and shoot ratio) of *Brachiaria brizantha*


TRATAMENTS		LDM	CDM	RDM	Root/shoot	K
Full sun	Non-inoculated	0.80 c	0.35 c	1.01 c	0.88 a	0.91 a
	BRM-32111	0.92 c	0.28 d	1.01 c	0.85 a	0.87 a
	BRM - 32113	1.32 b	0.42 b	1.08 b	0.63 c	0.64 c
	MIX	1.91 a	0.65 a	1.90 a	0.75 b	0.76 b
Shade	Non-inoculated	0.11 d	0.04 c	0.10 d	0.43 c	0.48 c
	BRM - 32111	0.29 b	0.14 b	0.37 b	0.90 c	0.96 b
	BRM - 32113	0.20 c	0.10 c	0.23 c	0.83 c	0.91 b
	MIX	0.44 a	0.24 a	0.77 a	1.16 a	1.21 a

1333 Means followed by different letters in each column and within each light intensity are
 1334 significantly different (P < 0.05, Duncan Test). BRM-3211 = *Pseudomonas fluorescens*; BRM-
 1335 3213 = *Burkholderia pyrrocinia*; MIX= BRM-3211 + BRM-3213.


1336

1337

1338 FIGURES



1339 **Figure 1** *Brachiaria brizantha* cv. Piatã under full-sun (a-d) and shade (e-h). 21 days after
 1340 inoculation (35 days after seedling emergence). Non-inoculated plant (a, e), inoculated with
 1341 *Pseudomonas fluorescens* (BRM-32111) (b, f), inoculated with *Burkholderia pyrrocinia*
 1342 (BRM-32113) (c, g) and co-inoculated with BRM-32111 + BRM-32113 (d, h).



1345 **Figure 2** Light intensity and plant growth-promoting rhizobacteria (PGPR) on plant height (H)
 1346 of *Brachiaria brizantha*. Columns with different letters are significantly different among PGPR
 1347 inoculation ($P < 0.05$, Duncan's test). Mean $\pm$ s.e. ($n = 5$). C = Non-inoculated; BRM-32111 =
 1348 *Pseudomonas fluorescens*; BRM-32113 = *Burkholderia pyrrocinia*; MIX = BRM-32111 +
 1349 BRM-32113.




Figure 3 Light intensity and plant growth-promoting rhizobacteria (PGPR) on specific leaf area (SLA) and leaf area ratio (LAR) of *Brachiaria brizantha*. Columns with different letters are significantly different among PGPR inoculation ($P < 0.05$, Duncan's test). Mean $\pm$ s.e. ($n = 5$). C = Non-inoculated; BRM-32111 = *Pseudomonas fluorescens*; BRM-32113 = *Burkholderia pyrrocinia*; MIX = BRM-32111 + BRM-32113.




Figure 4 Light intensity and plant growth-promoting rhizobacteria (PGPR) on leaf area (LA) and SPAD index (chlorophyll content) of *Brachiaria brizantha*. Columns with different letters are significantly different among PGPR inoculation ($P < 0.05$, Duncan's test). Mean $\pm$ s.e. ($n = 5$). C = Non-inoculated; BRM-32111 = *Pseudomonas fluorescens*; BRM-32113 = *Burkholderia pyrrocinia*; MIX = BRM-32111 + BRM-32113




Figure 5 Light intensity and plant growth-promoting rhizobacteria (PGPR) on total dry mass production (TDM) and relative growth rate (RGR) of *Brachiaria brizantha*. Columns with different letters are significantly different among PGPR inoculation ($P < 0.05$, Duncan's test). Mean $\pm$ s.e. ($n = 5$). C = Non-inoculated; BRM-32111 = *Pseudomonas fluorescens*; BRM-32113 = *Burkholderia pyrrocinia*; MIX= BRM-32111 + BRM-32113.




Figure 6 Light intensity and plant growth-promoting rhizobacteria (PGPR) on biomass allocation pattern of *Brachiaria brizantha*. Columns with different lower case letters are significantly different among treatments ($P < 0.05$, Duncan Test). Different upper-case letters within columns indicate significant differences among plant organs ($P < 0.05$, Duncan's test). Mean $\pm$ s.e. ($n = 5$). BRM-32111 = *Pseudomonas fluorescens*; BRM-32113 = *Burkholderia pyrrocinia*; MIX= BRM-32111 + BRM-32113.

1386 CONCLUSÕES GERAIS

O método de inoculação e a intensidade de luz foram fundamentais para determinar o potencial de *P. fluorescens* e *B. pyrrocinia* em promover o crescimento em *Brachiaria brizantha* cv. Piatã. A coinoculação dessas rizobactérias, pela rega do solo, modificou a anatomia (folha, colmo, raiz), incrementou as taxas fotossintéticas, eficiência instantânea do uso da água, índice Spad (clorofila), nitrato, proteínas, carboidratos totais, amido, N, P, Na, Mg e Fe, promovendo uma maior qualidade e produção de biomassa em *B. brizantha*. Também promoveram o crescimento em *B. brizantha* sob diferentes condições luminosas, incrementando em mais de 100% a produção de biomassa, em comparação as plantas controles com e sem adubo. Esse estudo evidencia o potencial biofertilizante da coinoculação de *P. fluorescens* e *B. pyrrocinia* em *B. brizantha* sob diferentes intensidades de luz, possibilitando seu uso como biofertilizante em sistemas convencionais e integrados, contribuindo para a obtenção de pastagens mais produtivas e com maior valor nutritivo.